

Review

Bio-Preservation of Meat and Fermented Meat Products by Lactic Acid Bacteria Strains and Their Antibacterial Metabolites

Shima Kaveh ¹, Seyed Mohammad Bagher Hashemi ^{2,*}, Elahe Abedi ², Mohammad Javad Amiri ^{3,*} and Francesca Laura Conte ⁴

¹ Faculty of Food Science and Technology, Gorgan University of Agricultural Sciences & Natural Resources, Gorgan 49189-43464, Iran; shima.kaveh_s97@gau.ac.ir

² Department of Food Science and Technology, College of Agriculture, Fasa University, Fasa 74616-86131, Iran; e.abedi@fasau.ac.ir

³ Department of Water Engineering, Faculty of Agriculture, Fasa University, Fasa 74616-86131, Iran

⁴ Department of Veterinary Sciences, University of Messina, Viale Giovanni Palatucci 13, 98168 Messina, Italy; francesca.conte@unime.it

* Correspondence: hashemi@fasau.ac.ir (S.M.B.H.); mj_amiri@fasau.ac.ir (M.J.A.)

Abstract: Meat and some meat products are highly perishable due to their high-water content, pH, and high content of nutrients. Therefore, spoilage control in these products is one of the critical challenges in the food industry. On the other hand, the increasing widespread awareness about the undesirable effects of synthetic preservatives has promoted the breakthrough of the use of natural compounds or bio-preservation technology. Bio-preservation implies the application of microorganisms or their metabolites to extend the shelf life of food products. In this regard, according to the ancient and safe use of fermentation by lactic acid bacteria (LAB), their application in the bio-preservation of meat and meat products is gaining more attention. Thus, more understanding of the potential of LAB and their metabolites in the control of pathogens in meat and meat products can create new horizons in the production of safe and functional products with long shelf life. So, this article aims to review the recent knowledge about the bio-preservation of meat and meat products by LAB and their metabolites. Also, their antibacterial mechanism and potential for use in hurdle technology are discussed. The outcome of this review literature shows the high potential of various LAB strains and their metabolites especially bacteriocins as bio-preservatives in meat and meat products for extending their shelf life. In this regard, their combined use with other novel technologies or natural antibacterial compounds as hurdle technology is a more effective method that can compete with synthetic preservatives.

Keywords: bio-preservation; fermentation; lactic acid bacteria; meat products; natural anti-microbial

Citation: Kaveh, S.; Hashemi, S.M.B.; Abedi, E.; Amiri, M.J.; Conte, F.L. Bio-Preservation of Meat and Fermented Meat Products by Lactic Acid Bacteria Strains and Their Antibacterial Metabolites.

Sustainability **2023**, *15*, 10154.

<https://doi.org/10.3390/su151310154>

Academic Editor: Đurđica Ačkar

Received: 12 May 2023

Revised: 13 June 2023

Accepted: 22 June 2023

Published: 26 June 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Meat and meat products are rich sources of nutrients for humans due to their high content of vitamin B groups, protein, essential amino acids, and minerals. Also, they provide a favorable environment for the growth of several microorganisms due to their ideal pH, nutrient factors, and high water activity [1]. The main bacteria involved in meat spoilage include the genera *Brochothrix*, *Enterobacter*, *Acinetobacter*, *Moraxella*, *Pseudomonas*, *Leuconostoc*, and *Proteus*, meanwhile some of them (i.e., *Enterobacter*, *Pseudomonas*) secrete biogenic amines, which might cause food safety issues [2]. Furthermore, meat and meat products may be contaminated by pathogenic microorganisms such as *Campylobacter jejuni*, *Clostridium botulinum*, *Escherichia coli*, *Bacillus cereus*, *Listeria monocytogenes*, *Clostridium perfringens*, *Salmonella* spp., *Yersinia enterocolitica*, and *Staphylococcus aureus* [3,4].

One of the main concern for the meat industry is the spoilage of fresh meat and meat products caused by microbial contamination [5]. The meat industry applies different

techniques to inhibit microbial growth and the production safe products with the suitable and desired shelf life [6]. Accordingly, the most common applied techniques include physical (e.g., drying, freezing, heat treatment, packaging, and curing) and especially chemical (e.g., use of synthetic preservative compounds) methods [7]. Nevertheless, chemical additives have many disadvantages such as the alteration of the nutritional and organoleptic properties of foods [8,9]. Also, the carcinogenicity and toxicity of many chemicals such as nitrates have been proven. Nitrates are the most common chemicals used in the meat industry for the inhibition of microbial growth, retardation of lipid oxidation, development of better flavor, taste, and aroma, and preserve the color of the meat. In fermented sausages, their conversion into nitrites by microbial nitrate reductases can inhibit the growth of spoilage bacteria such as *Clostridium* spp. [10]. It has been reported that their excess consumption can have dangerous effects on consumer health due to the formation of carcinogenic nitrosamines [11]. For example, nitrates can generate nitric oxide by nitrosation reactions, which can undergo a reaction with secondary amines and form N-nitrosamines [12,13]. So, the increasingly negative perceptions of synthetic preservative chemicals, the greater attention of consumers towards food quality, and increasing demand for high nutritional and synthetic chemical-free products has promoted the food industry to replace traditional preservation methods with green techniques, such as active packaging, modified atmosphere packaging, high hydrostatic pressure, pulsed electric fields, and bio-preservation [14–17]. In this field, bio-preservation is the most reliable and potent technique closely related to “from farm to fork” strategy. Bio-preservation is considered a method to extend the shelf life of food products using compounds derived from plants, animals, bacteria, or fungi [18]. However, most researchers focus on bio-preservation by using beneficial microorganisms and/or their antimicrobial compounds [19].

In this context, lactic acid bacteria (LAB) have attracted more attention than other bio-preservative microorganisms due to different reasons such as their encapsulation capability by extrusion during the production of the antimicrobial film [20] and their GRAS status approved by the U.S. Food and Drug Administration (FDA) as a preservative in some food [3,21]. On the other hand, increasing the demand for natural preservative methods in line with environmental protection led to an increase in researchers’ interest in finding efficient and sustainable preservative methods. In this regard, bio-preservation with LAB bacteria has no negative effects on consumers’ health and environment and have recently gained more attention as a useful and sustainable approach for the production of functional foods that lead to the sustainability of the consumers’ health [22,23]. So, the use of LAB and/or their metabolites, either alone or in combination with a low amount of natural or synthetic preservatives and moderate physicochemical treatments, may be an efficient solution to extend the shelf life and enhanced food safety (e.g., dairy products, fermented meat, and meat products) without negative effects on their nutritional quality [24]. Accordingly in the last two decades, intensive investigations have been focused on LAB and their antimicrobial metabolites to discover new LAB strains with food preservation potential to be used in sustainable preservative methods [25].

Hence, this review summarizes the current research on the bio-preservation of meat and meat products by LAB and their metabolites. Moreover, their mechanism of action and their application in hurdle technology and active packaging will be discussed.

2. Fermented Meat Products and Their Health-Beneficial Properties

In the past, different techniques were used and developed for the preservation of meat and meat products, starting with adding some ingredients, such as salt and sugar to reduce microorganisms without an exact understanding of their preservative mechanisms. But today, the use of microorganisms in terms of fermentation of meat and meat products is known as an effective preservation method [5]. Microbial enzymatic activities during fermentation leads to various physicochemical and microbial changes based on the meat components (natural or added components) [26]. Fermentation can occur by two pathways: (1) the use of natural microflora of meat or (2) the use of starter cultures such as lactic acid

bacteria and micrococci. Lactic acid bacteria breakdown the carbohydrates and micrococci reduce nitrates and nitrites to nitric oxide that leads to production of volatile and nonvolatile compounds and flavor and odor changes of the product [27]. Also, fermentation causes different health-beneficial properties in fermented meat products in comparison to non-fermented ones such as antioxidant, antimicrobial, anti-hypertensive, and antithrombotic [28,29]. Furthermore, some nutritional components are produced during fermentation that have the potential to prevent diabetes, cancers, and allergic sensitization [30]. More studies are needed to discover other health-beneficial potentials of these products and their exact mechanisms. Also, these products should be evaluated in terms of food safety.

3. A Brief Overview on LAB

LAB are part of the natural microbial flora of fermented meats and the intestinal microbiota of humans. These aerotolerant bacteria are mainly non-sporing, Gram-positive, Catalase-negative, and have either a spherical-shaped or rod-shaped cell (Figure 1) [31]. LAB are microaerophilic organisms and preferably require anaerobic conditions for growth. They play an important role in food fermentations; in fact, LAB can ferment carbohydrates to high amounts of lactic acid as the final product (homofermentative bacteria); in addition to lactic acid, heterofermentative bacteria produce acetic acid, carbon dioxide, and ethanol, as by-products [32]. These organisms are acidophilus with the optimum acidic pH values of 5.5–6.2, but few can tolerate pH as low as 3.0 [33]. LAB are Generally Regarded As Safe (GRAS) according to the FDA and the European Food Safety Authority (EFSA) that have granted many LAB species Qualified Presumption of Safety status (QPS) [25,34,35]. Lactic acid bacteria possess considerable bioactive properties such as cholesterol reduction and antimicrobial properties, which has led to an increased interest in their effective role as preservatives in innovative food preservation technology, much more than their application in traditional fermentation [36–38]. The antibacterial activity of LAB strains has been proven in different studies [39]. It has been reported that different LAB strains secrete various compounds that inhibit bacterial growth such as diacetyl, phenyl-lactate, organic acids, hydroxy fatty acid, hydroxy phenyl-lactate, hydrogen peroxide, propionate, and cyclic dipeptides. These bacteria also secrete biosurfactants, bacteriocins (i.e., acidophilin, lactacin, bifidocin, helveticin, plantarim, pediocin, bulgaricin, diplococin, and nisin), and bacteriocin-like inhibitory substances [40,41].




Figure 1. Schematic of properties of lactic acid bacteria.

3.1. LAB Strains Involved in Fermented Meat Products

The technological characterization of LAB strains involved in the fermentation process of meat is essential to select the best strain to be utilized as starter cultures [42]. The genera *Lactobacillus*, *Carnobacterium*, *Weissella*, *Pediococcus*, *Enterococcus*, and *Leuconostoc* are the main LAB that play crucial role in the fermentation [42,43]. The list of some of GRAS LAB that are most commonly used in bio-preservation of meat and meat products are mentioned in Table 1. Members of the genus *Lactobacillus* are usually the dominant species in most fermented meat products, but in some slightly acidified sausages, both *Enterococcus* and *Lactobacillus* are present in similar amounts [43]; nevertheless, *Lactobacillus plantarum* and *Lactobacillus curvatus* are the most common LAB species in fermented sausages [26]. It has been reported that in many fermented sausages, *Lactobacillus sakei* has the most adaptability due to a higher maximum growth rate, higher final cell density, and a shorter lag phase [44,45]. It should be noted that in Southern European sausages, the most and least common species are *Lactobacillus sakei* and *Pediococci* spp., respectively [46]. Also, molds, such as *Penicillium chrysogenum* and *Penicillium nalgiovense*, are commonly used for sausage ripening in Southern Europe [47]. In artisan sausages from Southern Europe, a strain of *Enterococcus faecium* grows increasingly during the early stages of fermentation, producing a bacteriocin [48]. It has been reported that yeast genera, especially *Debaryomyces hansenii*, can be found in fermented meat products with appropriate organoleptic characteristics [49,50]. In the next section, some data from the literatures on the bio-preservation of fermented meat products by application of starter LAB will be reviewed.

Table 1. Some of LAB species that are most commonly used in meat preservation.

Genus	Species	Genus	Species
<i>Lactobacillus</i>	<i>Lactobacillus delbrueckii</i>	<i>Lacticaseibacillus</i>	<i>Lacticaseibacillus paracasei</i>
	<i>Lactobacillus bulgaricus</i>		<i>Lacticaseibacillus rhamnosus</i>
	<i>Lactobacillus gallinarum</i>		<i>Lacticaseibacillus casei</i>
	<i>Lactobacillus gasseri</i>	<i>Pediococcus</i>	<i>Pediococcus acidilactici</i>
	<i>Lactobacillus lactis</i>		<i>Pediococcus pentosaceus</i>
	<i>Lactobacillus helveticus</i>		<i>Pediococcus parvulus</i>
	<i>Lactobacillus reuteri</i>	<i>Leuconostoc</i>	<i>Leuconostoc mesenteroides</i>
	<i>Lactobacillus acidophilus</i>		<i>Leuconostoc citreum</i>
	<i>Lactobacillus curvatus</i>		<i>Leuconostoc pseudomesenteroides</i>
	<i>Lactobacillus sakei</i>		<i>Leuconostoc carnosum</i>
<i>Lactobacillus salivarius</i>			
<i>Lactiplantibacillus</i>	<i>Lactiplantibacillus pentosus</i>	<i>Latilactobacillus</i>	<i>Latilactobacillus sakei</i>
	<i>Lactiplantibacillus plantarum</i>		<i>Latilactobacillus curvatus</i>
	<i>Lactiplantibacillus brevis</i>	<i>Limosilactobacillus</i>	<i>Limosilactobacillus fermentum</i>
	<i>Lactiplantibacillus casei</i>		<i>Limosilactobacillus reuteri</i>

3.2. Bio-Preservation of Meat and Meat Products by LAB and Their Metabolites

As mentioned above, bio-preservation strategies are based on the application of natural compounds derived from microorganisms, plants, or animals for extending the shelf life of food products [18]. But most studies circumscribe the bio-preservation concept to the application of microorganisms such as LAB or their metabolites to enhance food safety and extend the shelf life of food products. Generally, the most important approach is to use of microorganisms or their metabolites with antimicrobial activity against food spoilage bacteria and especially foodborne pathogens [25]. A desirable bio-preservation compound should only show antimicrobial activity against the targeted spoilage or pathogenic microorganisms and should not adversely affect the intestinal microbial flora of consumers [18].

LAB can be used directly as a functional ingredient in meat and meat products or as starters in fermentation processes. When directly applied, LAB can be added in freeze-dried or fresh cultures in different ways such as addition to fresh meat, meat batter formulation, or spraying on the surface of ready-to-eat meat products or fresh meat [25].

Traditionally, LAB have been widely used in fermentation processes, converting carbohydrates to lactic acid and producing biologically active compounds such as antibacterial and antifungal peptides, diacetyl, organic acids, and flavor precursors [51].

3.2.1. Bio-Preservation of Fermented Meat Products by Starter LAB

LAB play a key role in the production of fermented meat products resulting in texture and flavor improvement along with product preservation and finally the extension of shelf life. The strong buffering capacity and the low carbohydrate content of fresh meat lead to mild fermentation without variations in the organoleptic properties of food products. In naturally fermented meats, the addition of sugar leads to lactic acid production, pH decrease, and consequently protein denaturation by LAB activity [3]. The bio-preservation of meat and meat products by starter LAB has been reported in different studies as summarized in Table 2. For example, Nikodinoska et al. [52] evaluated the effect of *L. plantarum* PSC20 on microbial quality (*L. monocytogenes* and *Salmonella*) of Chorizo sausage. They reported that *L. plantarum* PSC20 caused a significant reduction in *L. monocytogenes* but did not have a significant effect on *Salmonella*. In another study, Lucumi-banguero et al. [53] revealed that *Lb. plantarum* and *Lb. sakei* have the maximum inhibition effect on *P. aureus* and the minimum effect on *S. typhimurium* and *S. marcescens*.

Table 2. The bio-preservative effect of LAB in meat and fermented meat products.

Product	Bio-Preservative LAB	Application Method	Targeted Microorganisms	Results	References
Vacuum-packaged fresh beef	<i>Lb. curvatus</i>	live culture and its bacteriocins, lactocin 705, and lactocin AL705	<i>L. monocytogenes</i> and <i>Br. thermosphacta</i>	Reduction in targeted microorganisms	[54]
Dry fermented sausage	<i>P. acidilactici</i>	as a starter culture	<i>L. monocytogenes</i>	Reduction in <i>L. monocytogenes</i> counts by 3.3 log CFU/g	[55]
Sliced fresh beef	<i>P. acidilactici</i> and <i>P. pentosaceus</i>	as a starter culture	<i>S. typhimurium</i> and <i>L. monocytogenes</i>	Reduction in <i>Enterobacteriaceae</i> , <i>Staphylococcus</i> , coliforms, <i>L. monocytogenes</i> and <i>S. typhimurium</i>	[56]
Vacuum packed raw beef	<i>L. curvatus</i>	as a starter culture	<i>B. thermosphacta</i> <i>Pseudomonas</i> sp.	No considerable bio-protective effect on <i>Pseudomonas</i> sp. Significant inhibition effect on <i>B. thermosphacta</i> . Controlling the growth of natural spoilage lactic acid bacteria in meat	[57]
Beef meatballs	Bacteriocin produced by <i>L. plantarum</i>	0.3% bacteriocin	<i>E. coli</i> and <i>S. typhimurium</i>	Reduction the total plate counts. Inhibition effect on <i>E. coli</i>	[58]
Fresh beef meat	Lactococcin BZ	various amounts of lactococcin BZ (200–2500 AU/mL)	Total aerobic, psychrotrophic and mesophilic bacteria, lactic acid bacteria, faecal coliforms,	Reduction in mesophilic, coliforms, psychrotrophic. 3.9 log reduction in LAB. 6 log reduction in <i>L. innocua</i> .	[59]

			total coliforms, and <i>L. innocua</i>		
Sucuk sauses	<i>L. plantarum</i> strains (S50, S51, S72, S74)	as starter cultures	<i>L. monocytogenes</i>	Reduction in the <i>L. monocytogenes</i> counts by 2.74 log CFU/g in the presence of <i>L. plantarum</i> S50	[60]
Chorizo sausage	<i>L. plantarum</i> PSC20	as protective cultures	<i>L. monocytogenes</i> and <i>Salmonella</i>	Reduction in <i>L. monocytogenes</i> No significant effect on <i>Salmonella</i>	[52]
Minced beef	<i>Lb. acidophilus</i>	as protective cultures	<i>Enterobacteriaceae</i> , coliform and <i>S. aureus</i>	Inhibition effect on <i>S. aureus</i>	[61]
Chorizo	<i>Lb. plantarum</i> and <i>Lb. sakei</i>	as protective cultures	<i>L. monocytogenes</i> , <i>E. coli</i> , <i>S. typhimurium</i> , <i>S. marcescens</i> , <i>S. aureus</i> , <i>P. aeruginosa</i> , and <i>P. vulgaris</i>	The most inhibition activity on <i>P. aureus</i> , and the least on <i>S. typhimurium</i> and <i>S. marcescens</i> .	[53]
Fresh red beef minced meat	<i>Lb. acidophilus</i> and <i>B. animalis</i>	400 AU/g Bac FL 31 bacteriocin	<i>S. aureus</i>	Reduction in the <i>S. aureus</i> by 0.89 and 0.55 CFU/g by <i>Lb. acidophilus</i> and <i>B. animalis</i> , respectively	[62]
Fermented sausage	<i>L. curvatus</i> 54M16	as starter cultures	<i>Enterobacteriaceae</i> and staphylococci	Reduction in Staphylococci and <i>Enterobacteriaceae</i>	[63]
Raw and roasted pork	<i>Lb. paracasei</i> F2I2	spraying with bacterial suspension	Staphylococci and coliforms	Reduction in Staphylococci and coliforms by more than 3.83 and 3.48 log CFU/g	[64]
Fermented pork meat sausage	<i>L. curvatus</i> 54M16	as cultures	<i>L. monocytogenes</i>	Reduction in <i>L. monocytogenes</i> by 0.51 log CFU/g	[65]

3.2.2. Bio-Preservation of Meat and Meat Products by Bacteriocins from LAB

As mentioned above, LAB produce antimicrobial compounds such as bacteriocins that have the potential to limit the growth of pathogen or spoilage microorganisms. It has been reported that more than half of almost all bacterial species can produce bacteriocins [66]. In this regard, Ren et al. [67] reported the noticeable antibacterial activity of *L. rhamnosus* sp. A5 against *E. coli*, *B. subtilis*, *Salmonella*, and *S. aureus*. They revealed that its purified bacteriocins have a pronounced inhibitory effect against *E. coli* and *B. subtilis*.

Bacteriocins are small bioactive peptides extracellularly released by Gram-negative and Gram-positive bacteria such as LAB [68]. In addition to extending the shelf life, bacteriocins have other benefits such as reducing the risk of transmission of pathogenic microorganisms and using a smaller amount in the application of synthetic preservatives [69]. However, bacteriocins are somewhat like antibiotics, but they have significant differences for many properties [70]. Bacteriocins are more active at wider pH ranges and are inherently more resistant to higher temperatures than antibiotics. Moreover, they have fast-acting antimicrobial mechanisms even at very low concentrations, so the development of resistant microbial strains in target bacteria is unlikely to be observed. In addition, the sensibility to proteases represents a limit as many Gram-negative bacteria produce proteases that will hydrolyze bacteriocins and thus inhibit the activity of bacteriocins [71]. Generally, bacteriocins are classified based on their molecular size, chemical structure, bacterial source, mechanism of action, and heat stability. Bacteriocins produced by Gram-positive LAB are divided into four groups (class I, class II, class III, and class IV).

Class I bacteriocins are thermo-stable, small (<5 kDa), ribosomally synthesized peptides with non-proteogenic thioether amino acids, lanthionine (Lan), and/or methyllanthionine (MeLan), so they are called lantibiotics [72].

Class II bacteriocins have molecular weight < 10 kDa. They are thermo-stable, hydrophobic, with 30–60 amino acids, contain a helical amphiphilic structure, which leads to the depolarization of the membrane of bacterial cell and consequently death of pathogens. These bacteriocins are called non-lantibiotics, as Lan or MeLan is absent [73].

Class III bacteriocins are heat-sensitive, large with molecular weight > 30 kDa [74]. This group have two subclasses, IIIa (bacteriolysin) and IIIb. The first subclass IIIa includes different bacteriocins such as lysostaphin and enterolysin that show their pathogenic effect by bacterial cell wall lysis. On the other hand, subclass IIIb such as helveticin M that act by dissipating the membrane and reducing the intracellular ATP concentration [75].

Class IV bacteriocins have complex structural moieties, so they are not considered as true bacteriocins and do not show antimicrobial potential. Some of colicins and microcins produced by *E. coli* belong to this group because of their antigenicity, microbial targets, and protein size [75]. The antibacterial effect of some bacteriocins from LAB are summarized in Table 3. For instance, de Azevedo et al. [76] evaluated the effect of bacteriocin-like inhibitory substances from *Pediococcus pentosaceus* in ready-to-eat pork ham. The results showed that in artificially contaminated ready-to-eat pork ham, bacteriocin-like inhibitors had considerable inhibition effect on the growth of *L. seeligeri* for 6 days. Chakchouk-Mtibaa et al. [77] investigated the effect of Bacteriocin BacFL31 from *Enterococcus faecium* on Turkey meat. They stated its considerable inhibition effect on *L. monocytogenes* and *Salmonella typhimurium*.

Table 3. Antibacterial effect of bacteriocins from LAB in meat and meat products.

Product	Bacteriocins	Results	References
Ready-to-eat pork ham	bacteriocin-like inhibitory substances from <i>P. pentosaceus</i>	Inhibition of <i>L. seeligeri</i> by 1.74 log CFU/g— Lower weigh loss	[76]
Chicken meat pieces inoculated	Sonorensin from <i>B. sonorensis</i>	Inhibition of <i>L. monocytogenes</i> and <i>S. aureus</i>	[78]
Raw ground Turkey meat	Bacteriocin BacFL31 by <i>E. faecium</i> FL31	Inhibition of <i>L. monocytogenes</i> and <i>S. typhimurium</i> by 3.2 log CFU/g and <i>S. aureus</i> by 2.2 log CFU/g	[77]
Vacuum-packaged beef frankfurters	Semi-purified bacteriocins from <i>L. curvatus</i> or <i>L. sakei</i>	Reducing pathogens to below detection level	[79]
Vacuum-sealed hot dogs	Bacteriocins from <i>L. curvatus</i> L442 and <i>L. lactis</i> subsp. <i>cremoris</i> ATCC 14365	Reduction in <i>L. monocytogenes</i> by greater than 2 log CFU/hot dog	[80]
Fresh beef	Lactococcin BZ bacteriocin from <i>L. lactis</i>	Reduction in mesophilic, psychrotrophic, and lactic acid bacteria by 4.87, 3.50, and 3.94 log, respectively, and 1.90×10^4 and 1.04×10^2 CFU/g reduction in coliform and fecal coliform bacteria, respectively	[59]
Beef meat	Bacteriocins from <i>L. crustorum</i> MN047	Significant reduction in <i>E.coli</i> and <i>S. aureus</i> by 4.3 and 4.5 log CFU/mL, respectively	[81]
Raw chicken breast	Bacteriocin XJS01 from <i>Ligilactobacillus salivarius</i>	47% reduction in <i>S. aureus</i>	[82]
Beef meat	Bacteriocin BM1300 from <i>L. crustorum</i> MN047	More antimicrobial effect against <i>E. coli</i> than <i>S. aureus</i> , 3.44 and 5.40 log CFU/mL, respectively	[83]

Fresh raw beef meat	Bacteriocin BM1122 from <i>L. crustorum</i> MN047	Inhibition effect on <i>S. aureus</i> and <i>E. coli</i> by 4.75 and 5.89 log CFU/mL	[84]
Fresh pork sausage	Bacteriocin from <i>L. curvatus</i> UFV-NPAC1	Reduction effect on <i>L. monocytogenes</i> as its counts ranged from 2 to 3.5 log CFU/g in control sample and ranged from 1.0 to 2.0 log CFU/g in treated sample through 10 days of storage	[85]
Chicken meat	Pediocin Ach/PA-1 from <i>P. pentosaceus</i> OZF	Reduction in <i>L. monocytogenes</i> counts by 3.8 log through 14 days of storage	[86]
Portuguese traditional fermented meat sausages	Pediocin PA-1 from <i>P. acidilactici</i> HA-6111-2	Reduction in <i>L. monocytogenes</i> to undetectable levels	[87]
Vacuum-packaged ready-to-eat meats	Bacteriocins from <i>L. curvatus</i> , <i>L. lactis</i> , <i>P. acidilactici</i> , <i>E. faecium</i> , and <i>E. thailandicus</i>	Reduction in <i>L. monocytogenes</i> by greater than 2 log	[88]

Synthesis and Mode of Action of Bacteriocins

The synthesis mechanism of bacteriocins can often be induced by population increase, stress conditions, nutrient shortage, cation surfactants, inhibitors, and the type of nitrogen, carbon, and phosphate sources in the media [89].

Bacteriocins are positively charged molecules with hydrophobic parts, and they interact electrostatically with phosphate groups (with negative charges) of cell membranes [14,90]. In other words, bacteriocins bind to receptors located on the cell wall of pathogen or spoilage bacteria, and then, bacteriocins kill the bacteria with various support mechanisms (Figure 2). Different mechanisms are involved in microorganisms' destruction by bacteriocin such as inhibition of nucleic acid synthesis, change in the cell translator mechanism, interference in the protein synthesis, and unbalanced functioning of the cytoplasmic membrane affecting cell permeability and energy use [91]. Moreover, lantibiotics inhibit target cells by creating pores on the membrane, causing disruption in the pH gradient or the transmembrane potential, leading to the leakage of cellular materials [14].




Figure 2. Modes of action of bacteriocins.

3.2.3. Bio-Preservation of Meat and Meat Products by Biosurfactants from LAB

As mentioned above, LAB produce different bioactive compounds including bacteriocins and bio-surfactants. In recent years, biosurfactants have reattracted much attention because of their considerable bioactive potential such as antibacterial, antiproliferative, and antioxidant activity [92,93]. Bio-surfactants are non-toxic, biodegradable, and amphiphilic compounds that are produced by various microorganisms such as LAB as secondary metabolites [94,95]. Biosurfactant are usually produce in two modes: cell-bound or excreted [96]. They have a diverse and complex chemical structure such as lipopolysaccharides, lipoproteins, and polysaccharide-protein complexes [97]. These compounds show distinct surface activity and can form micellar aggregates that lead to the reduction of the interfacial and surface tension [94,98,99]. Biosurfactants show appropriate activity at a wide range of pH and even under undesirable conditions such as salinities and high temperatures; so, they are suitable for food and pharmaceutical applications [100,101]. Their antimicrobial properties can play an effective role in the preservation of food products, including meat products. They exert their antimicrobial potential through various mechanisms such as: (1) The prevention of biofilm formation by the reduction of the bacterial interaction with the surface through alteration of the charge and wettability of the surface [94,102]; (2) Interference in the normal function of the microorganisms by interaction with their intracellular constituents [103]; (3) Destruction of the cell walls and membranes of microorganisms [104,105]. Different studies evaluated the antibacterial potential of biosurfactants from LAB on food pathogens; a summary of these studies is listed in Table 4.

Table 4. The antibacterial effect of biosurfactants from LAB on food pathogens with emphasis on meat and meat products.

Pathogens	Sources	Biosurfactant	Results	References
Raw ground goat meat		Biosurfactants from <i>L. paracasei</i> and <i>L. casei</i>	Reduction in the total aerobic counts, <i>E. coli</i> MTCC 118, and <i>P. aeruginosa</i> MTCC 1934	[106]
Fresh beef		Biosurfactants from <i>L. paracasei</i>	100% inhibition of <i>Bacillus</i> sp. BC1, <i>S. aureus</i> STP1, and <i>S. xylosus</i> STP2	[107]
-----		Biosurfactants from <i>L. paracasei</i> , <i>L. delbrueckii</i> , and <i>L. acidophilus</i>	18, 47.36, and 30.92% inhibition of <i>S. aureus</i> , respectively	[108]
-----		Biosurfactants from <i>L. paracasei</i>	100% inhibition of <i>E. coli</i> , <i>S. agalactiae</i> and <i>S. pyogenes</i>	[109]
-----		Biosurfactants from <i>L. paracasei</i> ssp. <i>Paracasei</i> A20	Reduction in <i>C. albicans</i> , <i>S. aureus</i> , and <i>S. Epidermidis</i>	[110]
-----		Biosurfactants from <i>S. thermophilus</i> A	Antimicrobial activity against <i>S. epidermidis</i> , <i>S. aureus</i> , <i>S. salivarius</i> , <i>R. dentocariosa</i> , <i>C. albicans</i> , and <i>C. tropicalis</i> at concentration of 20 and 40 g/L	[111]
-----		Biosurfactants from <i>L. lactis</i> 53	Antimicrobial activity against <i>S. epidermidis</i> , <i>S. aureus</i> , <i>S. salivarius</i> , <i>R. dentocariosa</i> , <i>C. albicans</i> , and <i>C. tropicalis</i> at concentration of 20 and 40 g/L	[112]

3.3. LAB or Their Metabolites as a Part of Hurdle Technology

Hurdle technology refers to the combination of different preservative factors such as water activity (a_w), temperature, redox potential (Eh), and novel preservative techniques, such as gas packaging, natural extracts, essential oils, and bacteriocins, to create more selective and efficient defensive systems to overcome pathogenic and spoilage microorganisms [69]. So, LAB and their metabolites can be used as a part of hurdles technology that act synergistically and inhibit food spoilage in combination with other preservative agents. In this way, less intensities of technological treatment and/or doses of preservative agents are required [113]. In the application of different antimicrobial agents, it is very important to select the best combination, so that desirable preservative effects are

achieved. In this regard, it has been reported that the addition of chelating agents makes the outer membranes of Gram-negative bacteria more permeable and sensitive to the hydrophobic peptides such as bacteriocins [114]. Also, freezing temperatures and modified atmosphere packaging (MAP) in combination with LAB and their metabolites can be used in hurdles technology approaches [115]. Different studies have evaluated the effect of different LAB in combination with various bio-preservatives on the microbial quality of meat and fermented meat products (summarized in Table 5).

Table 5. The antibacterial effect of LAB or their metabolites in combination with other preservative agents in meat and meat products.

Product	Bio-Preservative Agent	Results	References
Ground beef	<i>L. reuteri</i> or <i>L. plantarum</i> in combination with garlic extract	1.4 and 1.5 log reduction of <i>L. monocytogenes</i> by using <i>L. reuteri</i> or <i>L. plantarum</i> in combination with 1% of garlic extract	[116]
Beef sausage	Bacteriocinogenic <i>Enterococcus mundtii</i> and 0.0075% ascorbic acid, 3% NaCl, 0.02% NaNO ₂ , 0.75% glucose and 0.75% sucrose	>2 log cfu/g reduction of <i>L. monocytogenes</i>	[117]
Sliced beef	Bacteriocin from <i>C. maltaromaticum</i> combined with steam and chitosan	No synergistic effect 2 log reduction of <i>S. typhimurium</i> <i>E. coli</i> and <i>S. typhimurium</i>	[118]
Minced beef meat	<i>Mentha piperita</i> essential oil with semi-purified bacteriocin	Reduction in <i>Enterobacteriaceae</i>	[119]
Frozen ground beef patties	Bacteriocin-producing <i>L. curvatus</i> and <i>L. lactis</i> in combination with Na ₂ EDTA	1 log reduction of <i>E. coli</i>	[120]
Fresh chicken meat burger	<i>L. pseudomesenteroides</i> combined with MAP (50% CO ₂ and 50% O ₂)	Reduction in <i>L. monocytogenes</i> and <i>C. jejuni</i>	[115]
Fresh pork sausage	Combination of essential oils, nisin, nitrite, and organic acid salts, encapsulated	Reduction in <i>L. monocytogenes</i>	[121]
Alheira paste	<i>L. sakei</i> and <i>L. plantarum</i> , vacuum packed or packed under MAP (20% CO ₂ , 80% N ₂)	2 log reduction in <i>L. monocytogenes</i> by <i>L. sakei</i> . No significant differences between vacuum or MAP	[122]
Sliced lombo	Combination of Bacteriocin from <i>P. acidilactici</i> with HPP	Reduction in <i>L. innocua</i>	[123]
Goat meat emulsion	Combination of Pediocin from <i>P. pentosaceus</i> and <i>Murraya koenigii</i> berries extract	Reduction in <i>L. innocua</i>	[124]
Ready-to-eat pork ham	Bacteriocin-like inhibitory substances (BLIS) from <i>P. pentosaceus</i> and nisin	Inhibition of growth of <i>L. seeligeri</i>	[76]

3.3.1. LAB Application in Active Packaging

In addition to increasing demand for natural preservative agents, the demand for biodegradable and ecologically friendly packaging materials is increasing. On the other hand, the direct use of preservative agents in meat and meat products may reduce their bio-active properties such as their antimicrobial activity. So, in order to avoid this negative effect, the application of preservative agents as one of the components of active packaging is a suitable solution. Active packaging is a promising technology that actively modifies the internal environment of the food product package by interacting with the food over the storage time. Also, it is defined as an intelligent packaging system that alter and modify the environment inside the package and consequently the state of the food system in

order to improve the food quality and extend the shelf life of the product [125,126]. In this field, anti-microbial packaging is an efficient type of active packaging that have attracted much attention. Anti-microbial packaging has the ability to inhibit or kill pathogenic or spoilage food contamination microorganisms [127]. Antimicrobial agents can initially incorporate into the packaging materials and migrate into the food through partitioning and diffusion. Different approaches involve the incorporation of antimicrobial agents into food packaging, such as adding antimicrobial agents into the film formulation and adding them in the extruder when producing the film. Generally, different factors affect the properties of antimicrobial film such as the condition of casting process, the antimicrobial potential of the antimicrobial agent, physicochemical properties of film, mass transfer coefficient, and storage temperature [127,128]. In this regard, the application of preservative agents such as LAB and their metabolites in active packaging has various advantages such as controlled release and the need to lower the amount of these agents [113,129]. Various studies have investigated the preservative effect of LAB and their bacteriocins on the quality of meat and meat products (summarized in Table 6).

Table 6. The application of LAB or their metabolite in active packaging of meat and meat products.

Product	Bio-preservative agent	Results	References
Natural and artificial casings of meat product	Sakacin G from <i>L. curvatus</i>	Reduction in <i>L. innocua</i>	[130]
A pullulan film in ready-to-eat turkey breasts	Sakacin A from <i>Lactobacillus sakei</i> DSMZ 6333	Reduction in <i>L. monocytogenes</i>	[131]
An active polyvinylidene chloride film on fresh pork	Plantaricin from <i>Lactococcus plantarum</i> BM-1	Reduction in <i>L. monocytogenes</i>	[132]
A novel biocomposite film made of poly lactic acid and sawdust particles on raw sliced pork	Pediocin PA-1/AcH	Reduction in <i>L. monocytogenes</i>	[133]

3.4. Kinetics Models for Microbial Inactivation

Different kinetic models have been widely used for predicting the inactivation patterns of microorganisms. In this regard, the first-order kinetic mode is employed for log-linear survival curves, while the Weibull, biphasic, and log-logistic models are used for non-log-linear inactivation patterns. The first-order kinetic mode, the Weibull, biphasic, and log-logistic are expressed by the following equations, respectively [134]:

$$\log \frac{N_t}{N_0} = -\frac{t}{D_T} \quad (1)$$

$$\log \frac{N_t}{N_0} = -bt^n \quad (2)$$

$$\log N_t = \log N_0 + \log (k \times e^{-\alpha t} + (1 - k) \times e^{-\beta t}) \quad (3)$$

$$\log \frac{N_t}{N_0} = \frac{A}{1 + e^{\frac{4\sigma(\tau - \log t)}{A}}} + \frac{A}{1 + e^{\frac{4\sigma(\tau + 6)}{A}}} \quad (4)$$

where N_0 and N_t are the initial and surviving populations of bacteria at any time (CFU/g), t is time (min), D_T is defined as the time at which 90% of the bacterial population is inactivated, b is the inverse of the shape factor (1/min), n is the shape parameter (dimensionless), α and β are the inactivation kinetic rate constants (1/min), σ is the maximum inactivation rate (log (CFU/g)/log min), τ is the log time to attain the maximum inactivation rate (log min), A is the log increase in population. The statistical criteria are used to determine the goodness of fit of the kinetic models for describing the survival data.

4. Conclusions

The further negative perceptions related to the increase of synthetic preservative chemicals in meat products and the increased consumer focus on the relationship between the daily diet quality and health has led the food researchers to replace synthetic preservatives with natural compounds. Generally, an effective bio-preservation compound should only show antimicrobial activity against the targeted spoilage or pathogenic microorganisms and should not adversely affect the intestinal human microbial flora. In this regard, LAB and their metabolites have been widely studied due to their high preservative activity, which can potentially be used in the bio-preservation of meat and fermented meat products. In this paper, a general overview of LAB was given with reference to the species involved in fermentation of meat products. The genera of *Lactobacillus*, *Carnobacterium*, *Weissella*, *Pediococcus*, *Enterococcus*, and *Leuconostoc* are the leading LAB that play an essential role in fermentation. LAB produce various biological compounds such as antibacterial and antifungal peptides, diacetyl, organic acids, etc. In this field, bacteriocins are small bioactive peptides that are released by Gram-negative and Gram-positive bacteria such as LAB and induce a shelf-life extension of meat products. On the other hand, the combined use of LAB or their metabolites with other preservatives (nisin, natural extracts, essential oils, etc.) and novel techniques (MAP, HPP, active packaging, etc.) as hurdle technology, significantly increase their preservative effect. Moreover, the incorporation of LAB and their metabolite in active packaging is a more efficient method in their application as bio-preservatives in meat and meat products in comparison to direct application. As a result, these conclusions in the application of LAB in bio-preservation will pave the way for commercial use of LAB and their metabolites, especially bacteriocins, in the meat industry as natural preservatives to replace synthetic compounds. In addition, the combination use of these microbial bio-preservatives with other antibacterial compounds has an effective result on the shelf life and security of meat products. However, it is necessary to evaluate and characterize the novel bacteriocins and enhance suitable preservation techniques to avoid resistance.

Author Contributions: Conceptualization, S.M.B.H. and S.K.; supervision, E.A.; writing—original draft preparation, S.K.; writing—review and editing; M.J.A. and F.L.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data are available in the paper.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Bohrer, B.M. Nutrient Density and Nutritional Value of Meat Products and Non-Meat Foods High in Protein. *Trends Food Sci. Technol.* **2017**, *65*, 103–112.
2. Gao, X.; Li, C.; He, R.; Zhang, Y.; Wang, B.; Zhang, Z.-H.; Ho, C.-T. Research Advances on Biogenic Amines in Traditional Fermented Foods: Emphasis on Formation Mechanism, Detection and Control Methods. *Food Chem.* **2022**, *405*, 134911.
3. Favaro, L.; Todorov, S.D. Bacteriocinogenic LAB Strains for Fermented Meat Preservation: Perspectives, Challenges, and Limitations. *Probiotics Antimicrob. Proteins* **2017**, *9*, 444–458.
4. Laukkanen-Ninios, R.; Fredriksson-Ahomaa, M.; Korkeala, H. Enteropathogenic *Yersinia* in the Pork Production Chain: Challenges for Control. *Compr. Rev. Food Sci. Food Saf.* **2014**, *13*, 1165–1191.
5. Ashaolu, T.J.; Khalifa, I.; Mesak, M.A.; Lorenzo, J.M.; Farag, M.A. A Comprehensive Review of the Role of Microorganisms on Texture Change, Flavor and Biogenic Amines Formation in Fermented Meat with Their Action Mechanisms and Safety. *Crit. Rev. Food Sci. Nutr.* **2021**, *20*, 1–18.
6. Deak, T. Thermal Treatment. In *Food Safety Management*; Elsevier: Amsterdam, The Netherlands, 2014; pp. 423–442.
7. Buncic, S.; Nychas, G.-J.; Lee, M.R.F.; Koutsoumanis, K.; Hébraud, M.; Desvaux, M.; Chorianopoulos, N.; Bolton, D.; Blagojevic, B.; Antic, D. Microbial Pathogen Control in the Beef Chain: Recent Research Advances. *Meat Sci.* **2014**, *97*, 288–297.

8. Kaveh, S.; Mahoonak, A.S.; Ghorbani, M.; Jafari, S.M. Fenugreek Seed (*Trigonella Foenum Graecum*) Protein Hydrolysate Loaded in Nanosized Liposomes: Characteristic, Storage Stability, Controlled Release and Retention of Antioxidant Activity. *Ind. Crops Prod.* **2022**, *182*, 114908.
9. Radi, M.; Shadikhah, S.; Sayadi, M.; Kaveh, S.; Amiri, S.; Bagheri, F. Effect of *Thymus Vulgaris* Essential Oil-Loaded Nanostructured Lipid Carriers in Alginate-Based Edible Coating on the Postharvest Quality of Tangerine Fruit. *Food Bioprocess Technol.* **2023**, *16*, 185–198.
10. Duarte, M.; de Fátima Carrijo, K. Quantificação Do Teor de Nitrito de Sódio Residual Em Linguças Cozidas Tipo Calabresa Comercializadas No Sul Do Estado Do Rio de Janeiro, Brasil. *Enciclopédia Biosf.* **2014**, *10*, 1606–1615.
11. Eskandari, M.H.; Hosseinpour, S.; Mesbahi, G.; Shekarforoush, S. New Composite Nitrite-free and Low-nitrite Meat-curing Systems Using Natural Colorants. *Food Sci. Nutr.* **2013**, *1*, 392–401.
12. Bhattacharya, D.; Nanda, P.K.; Pateiro, M.; Lorenzo, J.M.; Dhar, P.; Das, A.K. Lactic Acid Bacteria and Bacteriocins: Novel Biotechnological Approach for Biopreservation of Meat and Meat Products. *Microorganisms* **2022**, *10*, 2058.
13. Flores, M.; Toldrá, F. Chemistry, Safety, and Regulatory Considerations in the Use of Nitrite and Nitrate from Natural Origin in Meat Products—Invited Review. *Meat Sci.* **2021**, *171*, 108272.
14. Hernández-Aquino, S.; Miranda-Romero, L.A.; Fujikawa, H.; Maldonado-Simán, E.D.E.J.; Alarcon-Zuniga, B. Antibacterial Activity of Lactic Acid Bacteria to Improve Shelf Life of Raw Meat. *Biocontrol Sci.* **2019**, *24*, 185–192.
15. Jafarpour, D.; Hashemi, S.M.B. Pure and Co-Fermentation of Quinoa Seeds by *Limosilactobacillus Fermentum* and *Lacticaeibacillus Rhamnosus*: Bioactive Content, Antidiabetic and Antioxidant Activities. *Fermentation* **2023**, *9*, 80.
16. Kaveh, S.; Gholamhosseinpour, A.; Mohammad, S.; Hashemi, B.; Jafarpour, D.; Castagnini, J.M. Original Article Recent Advances in Ultrasound Application in Fermented and Non-Fermented Dairy Products: Antibacterial and Bioactive Properties. *Int. J. Food Sci. Technol.* **2023**, *58*, 3591–3607. <https://doi.org/10.1111/ijfs.16457>.
17. Rezazadeh-Bari, M.; Najafi-Darmian, Y.; Alizadeh, M.; Amiri, S. Numerical Optimization of Probiotic Ayran Production Based on Whey Containing Transglutaminase and Aloe Vera Gel. *J. Food Sci. Technol.* **2019**, *56*, 3502–3512.
18. Pisoschi, A.M.; Pop, A.; Georgescu, C.; Turcuş, V.; Olah, N.K.; Mathe, E. An Overview of Natural Antimicrobials Role in Food. *Eur. J. Med. Chem.* **2018**, *143*, 922–935.
19. Singh, V.P. Recent Approaches in Food Bio-Preservation—a Review. *Open Vet. J.* **2018**, *8*, 104–111.
20. Radosavljević, M.; Lević, S.; Pejin, J.; Mojović, L.; Nedović, V. Encapsulation Technology of Lactic Acid Bacteria in Food Fermentation. In *Lactic Acid Bacteria in Food Biotechnology*; Elsevier: Amsterdam, The Netherlands, 2022; pp. 319–347.
21. Register, F. Nisin Preparation: Affirmation of GRAS Status as a Direct Human Food Ingredient. *Fed Regist.* **1988**, *54*, 11247–11251.
22. Qiu, Y.; Lei, P.; Zhang, Y.; Sha, Y.; Zhan, Y.; Xu, Z.; Li, S.; Xu, H.; Ouyang, P. Recent Advances in Bio-Based Multi-Products of Agricultural Jerusalem Artichoke Resources. *Biotechnol. Biofuels* **2018**, *11*, 1–15.
23. Venegas-Ortega, M.G.; Flores-Gallegos, A.C.; Martínez-Hernández, J.L.; Aguilar, C.N.; Nevárez-Moorillón, G. V Production of Bioactive Peptides from Lactic Acid Bacteria: A Sustainable Approach for Healthier Foods. *Compr. Rev. Food Sci. Food Saf.* **2019**, *18*, 1039–1051.
24. Ananou, S.; Maqueda, M.; Martínez-Bueno, M.; Valdivia, E. Biopreservation, an Ecological Approach to Improve the Safety and Shelf-Life of Foods. *Commun. Curr. Res. Educ. Top. Trends Appl. Microbiol.* **2007**, *1*, 475–487.
25. Barcenilla, C.; Ducic, M.; López, M.; Prieto, M.; Álvarez-Ordóñez, A. Application of Lactic Acid Bacteria for the Biopreservation of Meat Products: A Systematic Review. *Meat Sci.* **2022**, *183*, 108661.
26. Ojha, K.S.; Kerry, J.P.; Duffy, G.; Beresford, T.; Tiwari, B.K. Technological Advances for Enhancing Quality and Safety of Fermented Meat Products. *Trends Food Sci. Technol.* **2015**, *44*, 105–116.
27. Lorenzo, J.M.; Bedia, M.; Bañón, S. Relationship between Flavour Deterioration and the Volatile Compound Profile of Semi-Ripened Sausage. *Meat Sci.* **2013**, *93*, 614–620.
28. Ashaolu, T.J.; Reale, A. A Holistic Review on Euro-Asian Lactic Acid Bacteria Fermented Cereals and Vegetables. *Microorganisms* **2020**, *8*, 1176.
29. Xing, L.; Liu, R.; Cao, S.; Zhang, W.; Guanghong, Z. Meat Protein Based Bioactive Peptides and Their Potential Functional Activity: A Review. *Int. J. Food Sci. Technol.* **2019**, *54*, 1956–1966.
30. Ashaolu, T.J. Safety and Quality of Bacterially Fermented Functional Foods and Beverages: A Mini Review. *Food Qual. Saf.* **2020**, *4*, 123–127.
31. Straffella, S.; Simpson, D.J.; Yaghoubi Khanghahi, M.; De Angelis, M.; Gänzle, M.; Minervini, F.; Crecchio, C. Comparative Genomics and in Vitro Plant Growth Promotion and Biocontrol Traits of Lactic Acid Bacteria from the Wheat Rhizosphere. *Microorganisms* **2020**, *9*, 78.
32. Ribeiro, S.C.; Coelho, M.C.; Silva, C.C.G. A Rapid Screening Method to Evaluate Acidifying Activity by Lactic Acid Bacteria. *J. Microbiol. Methods* **2021**, *185*, 106227.
33. Khalid, K. An Overview of Lactic Acid Bacteria. *Int. J. Biosci* **2011**, *1*, 1–13.
34. BIOHAZ; Koutsoumanis, K.; Allende, A.; Alvarez-Ordóñez, A.; Bolton, D.; Bover-Cid, S.; Chemaly, M.; Davies, R.; De Cesare, A.; Hilbert, F. Scientific Opinion on the Update of the List of QPS-recommended Biological Agents Intentionally Added to Food or Feed as Notified to EFSA (2017–2019). *EFSA J.* **2020**, *18*, e05966.
35. Zheng, J.; Wittouck, S.; Salvetti, E.; Franz, C.M.A.P.; Harris, H.M.B.; Mattarelli, P.; O’toole, P.W.; Pot, B.; Vandamme, P.; Walter, J. A Taxonomic Note on the Genus *Lactobacillus*: Description of 23 Novel Genera, Emended Description of the Genus *Lactobacillus* Beijerinck 1901, and Union of *Lactobacillaceae* and *Leuconostocaceae*. *Int. J. Syst. Evol. Microbiol.* **2020**, *70*, 2782–2858.

36. Saranraj, P.; Naidu, M.A.; Sivasakthivelan, P. Lactic Acid Bacteria and Its Antimicrobial Properties: A Review. *Int. J. Pharm. Biol. Arch.* **2013**, *4*, 1124–1133.
37. Hashemi, S.M.B.; Abedi, E.; Kaveh, S.; Mousavifard, M. Hypocholesterolemic, Antidiabetic and Bioactive Properties of Ultrasound-Stimulated Exopolysaccharide Produced by *Lactiplantibacillus Plantarum* Strains. *Bioact. Carbohydr. Diet. Fibre* **2022**, *28*, 100334. <https://doi.org/10.1016/j.bcdf.2022.100334>.
38. Walhe, R.A.; Diwanay, S.S.; Patole, M.S.; Sayyed, R.Z.; Al-Shwaiman, H.A.; Alkhulaifi, M.M.; Elgorban, A.M.; Danish, S.; Datta, R. Cholesterol Reduction and Vitamin B12 Production Study on *Enterococcus Faecium* and *Lactobacillus Pentosus* Isolated from Yoghurt. *Sustainability* **2021**, *13*, 5853.
39. Al-Dhabi, N.A.; Esmail, G.A.; Valan Arasu, M. Co-Fermentation of Food Waste and Municipal Sludge from the Saudi Arabian Environment to Improve Lactic Acid Production by *Lactobacillus Rhamnosus* AW3 Isolated from Date Processing Waste. *Sustainability* **2020**, *12*, 6899.
40. Lee, N.-K.; Paik, H.-D. Prophylactic Effects of Probiotics on Respiratory Viruses Including COVID-19: A Review. *Food Sci. Biotechnol.* **2021**, *30*, 773–781.
41. Mei, J.; Ma, X.; Xie, J. Review on Natural Preservatives for Extending Fish Shelf Life. *Foods* **2019**, *8*, 490.
42. Hugo, C.J.; Hugo, A. Current Trends in Natural Preservatives for Fresh Sausage Products. *Trends Food Sci. Technol.* **2015**, *45*, 12–23.
43. Ammor, S.; Dufour, E.; Zagorec, M.; Chaillou, S.; Chevallier, I. Characterization and Selection of *Lactobacillus Sakei* Strains Isolated from Traditional Dry Sausage for Their Potential Use as Starter Cultures. *Food Microbiol.* **2005**, *22*, 529–538.
44. Hugas, M.; Garriga, M.; Aymerich, M.T. Functionality of Enterococci in Meat Products. *Int. J. Food Microbiol.* **2003**, *88*, 223–233.
45. Toldrá, F.; Hui, Y.H. Dry-Fermented Sausages and Ripened Meats: An Overview. In *Handbook of fermented meat and poultry*; Fidel Toldrá, Y. H. Hui, Iciar Astiasarán, Joseph G. Sebranek, R.T., Ed.; Wiley Online Library, 2014; pp. 1–6.
46. Franz, C.M.A.P.; Stiles, M.E.; Schleifer, K.H.; Holzapfel, W.H. Enterococci in Foods—A Conundrum for Food Safety. *Int. J. Food Microbiol.* **2003**, *88*, 105–122.
47. Leroy, F.; Verluyten, J.; De Vuyst, L. Functional Meat Starter Cultures for Improved Sausage Fermentation. *Int. J. Food Microbiol.* **2006**, *106*, 270–285.
48. Todorov, S.D.; Favaro, L.; Gibbs, P.; Vaz-Velho, M. *Enterococcus Faecium* Isolated from Lombo, a Portuguese Traditional Meat Product: Characterisation of Antibacterial Compounds and Factors Affecting Bacteriocin Production. *Benef. Microbes* **2012**, *3*, 319–330.
49. Andrade, M.J.; Córdoba, J.J.; Casado, E.M.; Córdoba, M.G.; Rodríguez, M. Effect of Selected Strains of *Debaryomyces Hansenii* on the Volatile Compound Production of Dry Fermented Sausage “Salchichón”. *Meat Sci.* **2010**, *85*, 256–264.
50. Cano-García, L.; Flores, M.; Belloch, C. Molecular Characterization and Aromatic Potential of *Debaryomyces Hansenii* Strains Isolated from Naturally Fermented Sausages. *Food Res. Int.* **2013**, *52*, 42–49.
51. Egan, K.; Field, D.; Rea, M.C.; Ross, R.P.; Hill, C.; Cotter, P.D. Bacteriocins: Novel Solutions to Age Old Spore-Related Problems? *Front. Microbiol.* **2016**, *7*, 461.
52. Nikodinoska, I.; Baffoni, L.; Di Gioia, D.; Manso, B.; García-Sánchez, L.; Melero, B.; Rovira, J. Protective Cultures against Food-borne Pathogens in a Nitrite Reduced Fermented Meat Product. *LWT* **2019**, *101*, 293–299.
53. Lucumi-banguero, R.S.; Ram, C. Potential Use of Lactic Acid Bacteria with Pathogen Inhibitory Capacity as a Biopreservative Agent for Chorizo. *Processes* **2021**, *9*, 1582.
54. Castellano, P.; Vignolo, G. Inhibition of *Listeria Innocua* and *Brochothrix Thermosphacta* in Vacuum-packaged Meat by Addition of Bacteriocinogenic *Lactobacillus Curvatus* CRL705 and Its Bacteriocins. *Lett. Appl. Microbiol.* **2006**, *43*, 194–199.
55. Coşansu Akdemir, S. Control of *Listeria Monocytogenes* by Bacteriocin-Producing *Pediococcus Aciditactici* 13 and Its Antimicrobial Substance in a Dry Fermented Sausage Sucuk and in Turkey Breast. *J. Food Nutr. Res.* **2010**, *49*, 206–214.
56. Olaoye, O.A.; Onilude, A.A. Investigation on the Potential Application of Biological Agents in the Extension of Shelf Life of Fresh Beef in Nigeria. *World J. Microbiol. Biotechnol.* **2010**, *26*, 1445–1454.
57. Castellano, P.; González, C.; Carduza, F.; Vignolo, G. Protective Action of *Lactobacillus Curvatus* CRL705 on Vacuum-Packaged Raw Beef. Effect on Sensory and Structural Characteristics. *Meat Sci.* **2010**, *85*, 394–401.
58. Arief, I.I.; Jenie, B.S.L.; Suryati, T.; Ayuningtyas, G.; Fuziawan, A. Antimicrobial Activity of Bacteriocin from Indigenous *Lactobacillus Plantarum* 2C12 and Its Application on Beef Meatball as Biopreservative. *J. Indones. Trop. Anim. Agric.* **2012**, *37*, 90–96.
59. Yıldırım, Z.; Yerlikaya, S.; Öncül, N.; Sakin, T. Inhibitory Effect of Lactococin BZ against *Listeria Innocua* and Indigenous Microbiota of Fresh Beef. *Food Technol. Biotechnol.* **2016**, *54*, 317–323.
60. Kamiloğlu, A.; Kaban, G.; Kaya, M. Effects of Autochthonous *Lactobacillus Plantarum* Strains on *Listeria Monocytogenes* in Sucuk during Ripening. *J. Food Saf.* **2019**, *39*, e12618.
61. Amine, R.A.E. Effect of Bio Preservation as a Modern Technology on Quality Aspects and Microbial Safety of Minced Beef Effect of Bio Preservation as a Modern Technology on Quality Aspects and Microbial Safety of Minced Beef. *Glob. J. Biotechnol. Biochem.* **2012**, *7*, 38–49.
62. Isa, J.K.; Razavi, S.H. The Use of *Lactobacillus Acidophilus* and *Bifidobacterium Animalis* Ssp. *Lactis* BB12, as Probiotics to Reduce the Risk of Food Poisoning in Minced Meat. *Appl. Food Biotechnol.* **2018**, *5*, 173–183.
63. Casaburi, A.; Di Martino, V.; Ferranti, P.; Picariello, L.; Villani, F. Technological Properties and Bacteriocins Production by *Lactobacillus Curvatus* 54M16 and Its Use as Starter Culture for Fermented Sausage Manufacture. *Food Control* **2016**, *59*, 31–45.
64. Calumba, K.F.A.; Obsioma, V.P.; Jawa, C.A.E.; Ong, D.G.S. Assessment of *Lactobacillus Paracasei* F2I2 as a Possible Biopreservative for Raw Pork. *Mindanao J. Sci. Technol.* **2019**, *17*, 1–17.

65. Giello, M.; La Stora, A.; De Filippis, F.; Ercolini, D.; Villani, F. Impact of *Lactobacillus Curvatus* 54M16 on Microbiota Composition and Growth of *Listeria Monocytogenes* in Fermented Sausages. *Food Microbiol.* **2018**, *72*, 1–15. <https://doi.org/10.1016/j.fm.2017.11.003>.
66. Cotter, P.D.; Hill, C.; Ross, R.P. Bacteriocins: Developing Innate Immunity for Food. *Nat. Rev. Microbiol.* **2005**, *3*, 777–788.
67. Ren, S.; Yuan, X.; Liu, F.; Fang, F.; Iqbal, H.M.N.; Zahran, S.A.; Bilal, M. Bacteriocin from *Lactocaseibacillus Rhamnosus* Sp. A5: Isolation, Purification, Characterization, and Antibacterial Evaluation for Sustainable Food Processing. *Sustainability* **2022**, *14*, 9571.
68. Martinez, F.A.C.; Balciunas, E.M.; Salgado, J.M.; González, J.M.D.; Converti, A.; de Souza Oliveira, R.P. Lactic Acid Properties, Applications and Production: A Review. *Trends Food Sci. Technol.* **2013**, *30*, 70–83.
69. Castellano, P.; Belfiore, C.; Fadda, S.; Vignolo, G. A Review of Bacteriocinogenic Lactic Acid Bacteria Used as Bioprotective Cultures in Fresh Meat Produced in Argentina. *Meat Sci.* **2008**, *79*, 483–499.
70. Balciunas, E.M.; Martinez, F.A.C.; Todorov, S.D.; de Melo Franco, B.D.G.; Converti, A.; de Souza Oliveira, R.P. Novel Biotechnological Applications of Bacteriocins: A Review. *Food Control* **2013**, *32*, 134–142.
71. O'Connor, P.M.; Ross, R.P.; Hill, C.; Cotter, P.D. Antimicrobial Antagonists against Food Pathogens: A Bacteriocin Perspective. *Curr. Opin. Food Sci.* **2015**, *2*, 51–57.
72. Zimina, M.; Babich, O.; Prosekov, A.; Sukhikh, S.; Ivanova, S.; Shevchenko, M.; Noskova, S. Overview of Global Trends in Classification, Methods of Preparation and Application of Bacteriocins. *Antibiotics* **2020**, *9*, 553.
73. Alvarez-Sieiro, P.; Montalbán-López, M.; Mu, D.; Kuipers, O.P. Bacteriocins of Lactic Acid Bacteria: Extending the Family. *Appl. Microbiol. Biotechnol.* **2016**, *100*, 2939–2951.
74. Sun, Z.; Wang, X.; Zhang, X.; Wu, H.; Zou, Y.; Li, P.; Sun, C.; Xu, W.; Liu, F.; Wang, D. Class III Bacteriocin Helveticin-M Causes Sublethal Damage on Target Cells through Impairment of Cell Wall and Membrane. *J. Ind. Microbiol. Biotechnol.* **2018**, *45*, 213–227.
75. Kumariya, R.; Garsa, A.K.; Rajput, Y.S.; Sood, S.K.; Akhtar, N.; Patel, S. Bacteriocins: Classification, Synthesis, Mechanism of Action and Resistance Development in Food Spoilage Causing Bacteria. *Microb. Pathog.* **2019**, *128*, 171–177.
76. de Azevedo, P.O.S.; Mendonça, C.M.N.; Seibert, L.; Domínguez, J.M.; Converti, A.; Gierus, M.; Oliveira, R.P.S. Bacteriocin-like Inhibitory Substance of *Pediococcus Pentosaceus* as a Biopreservative for *Listeria* Sp. Control in Ready-to-Eat Pork Ham. *Braz. J. Microbiol.* **2020**, *51*, 949–956.
77. Chakchouk-Mtibaa, A.; Smaoui, S.; Ktari, N.; Sellem, I.; Najah, S.; Karray-Rebai, I.; Mellouli, L. Biopreservative Efficacy of Bacteriocin BacFL31 in Raw Ground Turkey Meat in Terms of Microbiological, Physicochemical, and Sensory Qualities. *Biocontrol Sci.* **2017**, *22*, 67–77.
78. Chopra, L.; Singh, G.; Kumar Jena, K.; Sahoo, D.K. Sonorensin: A New Bacteriocin with Potential of an Anti-Biofilm Agent and a Food Biopreservative. *Sci. Rep.* **2015**, *5*, 1–13.
79. Castellano, P.; Peña, N.; Ibarreche, M.P.; Carduza, F.; Soteras, T.; Vignolo, G. Antilisterial Efficacy of *Lactobacillus* Bacteriocins and Organic Acids on Frankfurters. Impact on Sensory Characteristics. *J. Food Sci. Technol.* **2018**, *55*, 689–697.
80. Ünlü, G.; Nielsen, B.; Ionita, C. Inhibition of *Listeria Monocytogenes* in Hot Dogs by Surface Application of Freeze-Dried Bacteriocin-Containing Powders from Lactic Acid Bacteria. *Probiotics Antimicrob. Proteins* **2016**, *8*, 102–110.
81. Yan, H.; Lu, Y.; Li, X.; Yi, Y.; Wang, X.; Shan, Y.; Liu, B.; Zhou, Y.; Lü, X. Action Mode of Bacteriocin BM1829 against *Escherichia Coli* and *Staphylococcus Aureus*. *Food Biosci.* **2021**, *39*, 100794.
82. Li, H.-W.; Xiang, Y.-Z.; Zhang, M.; Jiang, Y.-H.; Zhang, Y.; Liu, Y.-Y.; Lin, L.-B.; Zhang, Q.-L. A Novel Bacteriocin from *Lactobacillus Salivarius* against *Staphylococcus Aureus*: Isolation, Purification, Identification, Antibacterial and Antibiofilm Activity. *LWT* **2021**, *140*, 110826.
83. Lu, Y.; Aizhan, R.; Yan, H.; Li, X.; Wang, X.; Yi, Y.; Shan, Y.; Liu, B.; Zhou, Y.; Lü, X. Characterization, Modes of Action, and Application of a Novel Broad-Spectrum Bacteriocin BM1300 Produced by *Lactobacillus Crustorum* MN047. *Braz. J. Microbiol.* **2020**, *51*, 2033–2048.
84. Lu, Y.; Yan, H.; Li, X.; Gu, Y.; Wang, X.; Yi, Y.; Shan, Y.; Liu, B.; Zhou, Y.; Lü, X. Physicochemical Properties and Mode of Action of a Novel Bacteriocin BM1122 with Broad Antibacterial Spectrum Produced by *Lactobacillus Crustorum* MN047. *J. Food Sci.* **2020**, *85*, 1523–1535.
85. de Castilho, N.P.A.; Todorov, S.D.; Oliveira, L.L.; dos Santos Bersot, L.; Nero, L.A. Inhibition of *Listeria Monocytogenes* in Fresh Sausage by Bacteriocinogenic *Lactobacillus Curvatus* UFV-NPAC1 and Its Semi-Purified Bacteriocin. *LWT* **2020**, *118*, 108757.
86. Kiran, F.; Osmanağaoğlu, Ö. Inhibition of *Listeria Monocytogenes* in Chicken Meat by *Pediococcus* AcH/PA-1 Produced by *Pediococcus Pentosaceus* OZF. *Agro Food Ind. Hi. Tech.* **2014**, *25*, 66–69.
87. Maciel, C.; Komora, N.; Ferreira, V.; Saraiva, J.; Castro, S.M.; Teixeira, P. High Hydrostatic Pressure and *Pediocin* PA-1 as a Synergistic System to *Listeria Monocytogenes* Inactivation in Fermented Meat Sausage. In Proceedings of the Congress of Microbiology and Biotechnology; Porto, Portugal, 2017.
88. Vijayakumar, P.P.; Muriana, P.M. Inhibition of *Listeria Monocytogenes* on Ready-to-Eat Meats Using Bacteriocin Mixtures Based on Mode-of-Action. *Foods* **2017**, *6*, 22.
89. Riley, M.A. Bacteriocin-Mediated Competitive Interactions of Bacterial Populations and Communities. In *Prokaryotic antimicrobial peptides: from genes to applications*; Drider, D., Rebuffat, S., Ed.; Springer, 2011; pp. 13–26.
90. Cleveland, J.; Montville, T.J.; Nes, I.F.; Chikindas, M.L. Bacteriocins: Safe, Natural Antimicrobials for Food Preservation. *Int. J. Food Microbiol.* **2001**, *71*, 1–20.
91. Lahiri, D.; Nag, M.; Sarkar, T.; Ray, R.R.; Shariati, M.A.; Rebezov, M.; Bangar, S.P.; Lorenzo, J.M.; Domínguez, R. Lactic Acid Bacteria (LAB): Autochthonous and Probiotic Microbes for Meat Preservation and Fortification. *Foods* **2022**, *11*, 2792.

92. Saravanakumari, P.; Mani, K. Structural Characterization of a Novel Xylo lipid Biosurfactant from *Lactococcus Lactis* and Analysis of Antibacterial Activity against Multi-Drug Resistant Pathogens. *Bioresour. Technol.* **2010**, *101*, 8851–8854.
93. Merghni, A.; Dallel, I.; Noumi, E.; Kadmi, Y.; Hentati, H.; Tobji, S.; Amor, A.B.; Mastouri, M. Antioxidant and Antiproliferative Potential of Biosurfactants Isolated from *Lactobacillus Casei* and Their Anti-Biofilm Effect in Oral *Staphylococcus Aureus* Strains. *Microb. Pathog.* **2017**, *104*, 84–89.
94. Banat, I.M.; Franzetti, A.; Gandolfi, I.; Bestetti, G.; Martinotti, M.G.; Fracchia, L.; Smyth, T.J.; Marchant, R. Microbial Biosurfactants Production, Applications and Future Potential. *Appl. Microbiol. Biotechnol.* **2010**, *87*, 427–444.
95. Jahan, R.; Bodratti, A.M.; Tsianou, M.; Alexandridis, P. Biosurfactants, Natural Alternatives to Synthetic Surfactants: Physicochemical Properties and Applications. *Adv. Colloid Interface Sci.* **2020**, *275*, 102061.
96. Mouafo, H.T.; Sokamte, A.T.; Mbawala, A.; Ndjouenkeu, R.; Devappa, S. Biosurfactants from Lactic Acid Bacteria: A Critical Review on Production, Extraction, Structural Characterization and Food Application. *Food Biosci.* **2022**, *46*, 101598.
97. Rodríguez, N.; Salgado, J.M.; Cortés, S.; Domínguez, J.M. Alternatives for Biosurfactants and Bacteriocins Extraction from *Lactococcus Lactis* Cultures Produced under Different PH Conditions. *Lett. Appl. Microbiol.* **2010**, *51*, 226–233.
98. Sharma, D.; Singh Saharan, B. Simultaneous Production of Biosurfactants and Bacteriocins by Probiotic *Lactobacillus Casei* MRTL3. *Int. J. Microbiol.* **2014**, *2014*, 698713.
99. Saharan, B.S.; Sahu, R.K.; Sharma, D. A Review on Biosurfactants: Fermentation, Current Developments and Perspectives. *Genet. Eng. Biotechnol. J.* **2011**, *2011*, 1–14.
100. Muthusamy, K.; Gopalakrishnan, S.; Ravi, T.K.; Sivachidambaram, P. Biosurfactants: Properties, Commercial Production and Application. *Curr. Sci.* **2008**, *94*, 736–747.
101. Henkel, M.; Müller, M.M.; Kügler, J.H.; Lovaglio, R.B.; Contiero, J.; Syldatk, C.; Hausmann, R. Rhamnolipids as Biosurfactants from Renewable Resources: Concepts for next-Generation Rhamnolipid Production. *Process Biochem.* **2012**, *47*, 1207–1219.
102. Ashraf, A.; Ahmed, A.A.; Fatma, I.; Zeinab, A.M. Characterization and Bioactivities of *Lactobacillus Plantarum* and *Pediococcus Acidilactici* Isolated from Meat and Meat Products. *Nat. Sci.* **2019**, *17*, 187–193. <https://doi.org/10.7537/marsnsj171219.26>.
103. Inès, M.; Dhouha, G. Lipopeptide Surfactants: Production, Recovery and Pore Forming Capacity. *Peptides* **2015**, *71*, 100–112.
104. Sambanthamoorthy, K.; Feng, X.; Patel, R.; Patel, S.; Paranavitana, C. Antimicrobial and Antibiofilm Potential of Biosurfactants Isolated from *Lactobacilli* against Multi-Drug-Resistant Pathogens. *BMC Microbiol.* **2014**, *14*, 1–9.
105. Hippolyte, M.T.; Augustin, M.; Hervé, T.M.; Robert, N.; Devappa, S. Application of Response Surface Methodology to Improve the Production of Antimicrobial Biosurfactants by *Lactobacillus Paracasei* Subsp. *Tolerans* N2 Using Sugar Cane Molasses as Substrate. *Bioresour. Bioprocess.* **2018**, *5*, 1–16.
106. Mouafo, H.T.; Mbawala, A.; Tanaji, K.; Somashekar, D.; Ndjouenkeu, R. Improvement of the Shelf Life of Raw Ground Goat Meat by Using Biosurfactants Produced by *Lactobacilli* Strains as Biopreservatives. *LWT* **2020**, *133*, 110071.
107. Mouafo, H.T.; Baomog, A.M.B.; Adjele, J.J.B.; Sokamte, A.T.; Mbawala, A.; Ndjouenkeu, R. Microbial Profile of Fresh Beef Sold in the Markets of Ngaoundéré, Cameroon, and Antiadhesive Activity of a Biosurfactant against Selected Bacterial Pathogens. *J. Food Qual.* **2020**, *2020*, 5989428.
108. Gomaa, E.Z. Antimicrobial and Anti-Adhesive Properties of Biosurfactant Produced by *Lactobacilli* Isolates, Biofilm Formation and Aggregation Ability. *J. Gen. Appl. Microbiol.* **2013**, *59*, 425–436.
109. Gudina, E.J.; Teixeira, J.A.; Rodrigues, L.R.; Gudi, E.J. Biointerfaces Isolation and Functional Characterization of a Biosurfactant Produced by *Lactobacillus Paracasei*. *Colloids Surf. B Biointerfaces* **2010**, *76*, 298–304. <https://doi.org/10.1016/j.colsurfb.2009.11.008>.
110. Gudiña, E.J.; Rocha, V.; Teixeira, J.A.; Rodrigues, L.R. Antimicrobial and Antiadhesive Properties of a Biosurfactant Isolated from *Lactobacillus Paracasei* Ssp. *Paracasei* A20. *Lett. Appl. Microbiol.* **2010**, *50*, 419–424.
111. Rodrigues, L.R.; Teixeira, J.A.; Van Der Mei, H.C.; Oliveira, R. Isolation and Partial Characterization of a Biosurfactant Produced by *Streptococcus Thermophilus* A. *Colloids Surf. B Biointerfaces* **2006**, *53*, 105–112. <https://doi.org/10.1016/j.colsurfb.2006.08.009>.
112. Rodrigues, R.; Teixeira, A.; Van Der Mei, H.C. Physicochemical and Functional Characterization of a Biosurfactant Produced by *Lactococcus Lactis* 53. *Colloids Surf. B Biointerfaces* **2006**, *49*, 79–86. <https://doi.org/10.1016/j.colsurfb.2006.03.003>.
113. Gálvez, A.; Abriouel, H.; López, R.L.; Omar, N. Ben Bacteriocin-Based Strategies for Food Biopreservation. *Int. J. Food Microbiol.* **2007**, *120*, 51–70.
114. Belfiore, C.; Castellano, P.; Vignolo, G. Reduction of *Escherichia Coli* Population Following Treatment with Bacteriocins from Lactic Acid Bacteria and Chelators. *Food Microbiol.* **2007**, *24*, 223–229.
115. Melero, B.; Diez, A.M.; Rajkovic, A.; Jaime, I.; Rovira, J. Behaviour of Non-Stressed and Stressed *Listeria Monocytogenes* and *Campylobacter Jejuni* Cells on Fresh Chicken Burger Meat Packaged under Modified Atmosphere and Inoculated with Protective Culture. *Int. J. Food Microbiol.* **2012**, *158*, 107–112.
116. Khalili Sadaghiani, S.; Aliakbarlu, J.; Tajik, H.; Mahmoudian, A. Anti-listeria Activity and Shelf Life Extension Effects of *Lactobacillus* along with Garlic Extract in Ground Beef. *J. Food Saf.* **2019**, *39*, e12709.
117. Orihuel, A.; Bonacina, J.; Vildoza, M.J.; Bru, E.; Vignolo, G.; Saavedra, L.; Fadda, S. Biocontrol of *Listeria Monocytogenes* in a Meat Model Using a Combination of a Bacteriocinogenic Strain with Curing Additives. *Food Res. Int.* **2018**, *107*, 289–296.
118. Hu, Z.Y.; Balay, D.; Hu, Y.; McMullen, L.M.; Gänzel, M.G. Effect of Chitosan, and Bacteriocin-Producing *Carnobacterium Maltaromaticum* on Survival of *Escherichia Coli* and *Salmonella Typhimurium* on Beef. *Int. J. Food Microbiol.* **2019**, *290*, 68–75.
119. Smaoui, S.; Hsouna, A.B.; Lahmar, A.; Ennouri, K.; Mtibaa-Chakchouk, A.; Sellem, I.; Najah, S.; Bouaziz, M.; Mellouli, L. Bio-Preservative Effect of the Essential Oil of the Endemic *Mentha Piperita* Used Alone and in Combination with BacTN635 in Stored Minced Beef Meat. *Meat Sci.* **2016**, *117*, 196–204.

120. Castellano, P.; Belfiore, C.; Vignolo, G. Combination of Bioprotective Cultures with EDTA to Reduce Escherichia Coli O157: H7 in Frozen Ground-Beef Patties. *Food Control* **2011**, *22*, 1461–1465.
121. Ghabraie, M.; Vu, K.D.; Huq, T.; Khan, A.; Lacroix, M. Antilisterial Effects of Antibacterial Formulations Containing Essential Oils, Nisin, Nitrite and Organic Acid Salts in a Sausage Model. *J. Food Sci. Technol.* **2016**, *53*, 2625–2633.
122. Vaz-Velho, M.; Jácomea, S.; Noronhab, L.; Todorovc, S.; Fonseca, S.; Pinheiro, R.; Moraisb, A.; Silvab, J.; Teixeira, P. Comparison of Antilisterial Effects of Two Strains of Lactic Acid Bacteria during Processing and Storage of a Portuguese Salami-like Product “Alheira”. *Chem. Eng.* **2013**, *32*, 1807–1812.
123. Castro, S.M.; Silva, J.; Casquete, R.; Queirós, R.; Saraiva, J.A.; Teixeira, P. Combined Effect of Pediocin BacHA-6111-2 and High Hydrostatic Pressure to Control Listeria Innocua in Fermented Meat Sausage. *Int. Food Res. J.* **2018**, *25*, 553–560.
124. Kumar, Y.; Kaur, K.; Shahi, A.K.; Kairam, N.; Tyagi, S.K. Antilisterial, Antimicrobial and Antioxidant Effects of Pediocin and *Murraya Koenigii* Berry Extract in Refrigerated Goat Meat Emulsion. *LWT-Food Sci. Technol.* **2017**, *79*, 135–144.
125. Balasubramanian, A.; Rosenberg, L.E.; Yam, K.; Chikindas, M.L. Antimicrobial Packaging: Potential vs. Reality—A Review. *J. Appl. Packag. Res.* **2009**, *3*, 193–221.
126. Suppakul, P.; Miltz, J.; Sonneveld, K.; Bigger, S.W. Active Packaging Technologies with an Emphasis on Antimicrobial Packaging and Its Applications. *J. Food Sci.* **2003**, *68*, 408–420.
127. Deshmukh, P.V.; Thorat, P.R. Bacteriocins: A New Trend in Antimicrobial Food Packaging. *Int. J. Adv. Res. Eng. Appl. Sci.* **2013**, *2*, 1–12.
128. Fu, Y.; Dudley, E.G. Antimicrobial-coated Films as Food Packaging: A Review. *Compr. Rev. Food Sci. Food Saf.* **2021**, *20*, 3404–3437.
129. Kristo, E.; Koutsoumanis, K.P.; Biliaderis, C.G. Thermal, Mechanical and Water Vapor Barrier Properties of Sodium Caseinate Films Containing Antimicrobials and Their Inhibitory Action on *Listeria Monocytogenes*. *Food Hydrocoll.* **2008**, *22*, 373–386.
130. Rivas, F.P.; Cayre, M.E.; Campos, C.A.; Castro, M.P. Natural and Artificial Casings as Bacteriocin Carriers for the Biopreservation of Meats Products. *J. Food Saf.* **2018**, *38*, e12419.
131. Trinetta, V.; Floros, J.D.; Cutter, C.N. Sakacin A-containing Pullulan Film: An Active Packaging System to Control Epidemic Clones of *Listeria Monocytogenes* in Ready-to-eat Foods. *J. Food Saf.* **2010**, *30*, 366–381.
132. Xie, Y.; Zhang, M.; Gao, X.; Shao, Y.; Liu, H.; Jin, J.; Yang, W.; Zhang, H. Development and Antimicrobial Application of Plantaricin BM-1 Incorporating a PVDC Film on Fresh Pork Meat during Cold Storage. *J. Appl. Microbiol.* **2018**, *125*, 1108–1116.
133. Woraprayote, W.; Kingcha, Y.; Amonphanpokin, P.; Krueenate, J.; Zendo, T.; Sonomoto, K.; Benjakul, S.; Visessanguan, W. Anti-*Listeria* Activity of Poly (Lactic Acid)/Sawdust Particle Biocomposite Film Impregnated with Pediocin PA-1/AcH and Its Use in Raw Sliced Pork. *Int. J. Food Microbiol.* **2013**, *167*, 229–235.
134. Esua, O.J.; Sun, D.-W.; Ajani, C.K.; Cheng, J.-H.; Keener, K.M. Modelling of inactivation kinetics of *Escherichia coli* and *Listeria monocytogenes* on grass carp treated by combining ultrasound with plasma functionalized buffer. *Ultrason. Sonochem.* **2022**, *88*, 106086.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.