


Review Article

Pharmacological Aspects and Biological Effects of Cannabigerol and Its Synthetic Derivatives

Fabrizio Calapai,¹ Luigi Cardia,² Emanuela Esposito,^{1,3} Ilaria Ammendolia,⁴ Cristina Mondello,⁴ Roberto Lo Giudice,⁵ Sebastiano Gangemi,⁵ Gioacchino Calapai ,⁴ and Carmen Mannucci⁴

¹Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Messina 98168, Italy

²Department of Human Pathology in Adulthood and Childhood “G. Barresi”, University Hospital “G. Martino” of Messina, Via Consolare Valeria 1, Messina 98123, Italy

³Genetics and Pharmacogenetics Unit, Policlinico Universitario “G. Martino”, University of Messina, Messina 98125, Italy

⁴Department of Biomedical and Dental Sciences and Morphofunctional Imaging, University of Messina, Messina 98125, Italy

⁵Department of Clinical and Experimental Medicine, Unit and School of Allergy and Clinical Immunology, University of Messina, Messina 98125, Italy

Correspondence should be addressed to Gioacchino Calapai; gcalapai@unime.it

Received 7 January 2022; Revised 28 July 2022; Accepted 5 August 2022; Published 8 November 2022

Academic Editor: Orazio Tagliatalata Scafati

Copyright © 2022 Fabrizio Calapai et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cannabigerol (CBG) is a cannabinoid from the plant *Cannabis sativa* that lacks psychotomimetic effects. Its precursor is the acidic form, cannabigerolic acid (CBGA), which is, in turn, a biosynthetic precursor of the compounds cannabidiol (CBD) and Δ^9 -tetrahydrocannabinol (THC). CBGA decarboxylation leads to the formation of neutral cannabinoid CBG, through a chemical reaction catalyzed by heat. On the basis of the growing interest in CBG and with the aim of highlighting scientific information on this phytocannabinoid, we focused the content of this article on its pharmacokinetic and pharmacodynamic characteristics and on its principal pharmacological effects. CBG is metabolized in the liver by the enzyme CYP2J2 to produce hydroxyl and di-oxygenated products. CBG is considered a partial agonist at the CB1 receptor (R) and CB2R, as well as a regulator of endocannabinoid signaling. Potential pharmacological targets for CBG include transient receptor potential (TRP) channels, cyclooxygenase (COX-1 and COX-2) enzymes, cannabinoid, 5-HT1A, and alpha-2 receptors. Pre-clinical findings show that CBG reduces intraocular pressure, possesses antioxidant, anti-inflammatory, and anti-tumoral activities, and has anti-anxiety, neuroprotective, dermatological, and appetite-stimulating effects. Several findings suggest that research on CBG deserves to be deepened, as it could be used, alone or in association, for novel therapeutic approaches for several disorders.

1. Introduction

The increased therapeutic potential of *Cannabis sativa* L. (fam. Cannabaceae; *C. sativa*) and the pharmacology of its chemical constituents need a more in-depth understanding in terms of its other constituents, rather than the more known phytocannabinoids Δ^9 -tetrahydrocannabinol (THC) and cannabidiol (CBD) (Figure 1). After the isolation of

THC, the main psychoactive constituent of *C. sativa* [1, 2], over 100 phytocannabinoids have been found in this plant, one of these being cannabigerol (CBG) [3].

A few years after the identification of THC (Figure 1), *in vivo* assays showed that the phytocannabinoid CBG is not psychotomimetic like the more well-known phytocannabinoid THC [4, 5]. However, it has been neglected and shadowed by THC for years because of its




FIGURE 1: Major phytocannabinoids.

lower concentration and, paradoxically, probably for its lack of psychotomimetic activity. CBG is the terpenophenolic phytocannabinoid precursor, in the plant *C. sativa*, of both THC and CBD [6]. CBG is not present only in this plant; beyond its discovery in *C. sativa*, it has also been found in the phytochemical profile of an extract from *Helichrysum umbraculigerum*, considered to be the most abundant source of CBG [7] (Figure 1).

In the plant, the cannabinoids THC and CBD are synthesized in an acidic (carboxylated) form, namely, cannabigerolic acid (CBGA) (Figure 1). CBGA undergoes a decarboxylation process depending on different factors, in particular, the speed of this process is related to the rise in temperature (Figure 1). Decarboxylation causing CBG formation is due to a simple chemical reaction catalyzed by heat, a conversion taking place at room temperature, but in a much slower way [8]. This intermediate cannabinoid is not present at significant concentrations in *Cannabis*, commonly occurring as a minor compound in terms of proportions [9, 10]. As happens for THC and CBD, the plant primarily synthesizes CBGA; this acidic form of the cannabinoid is unstable from the thermic point of view and can be decarboxylated when exposed to light or heat via smoking, baking, or refluxing [11]. In this article, we highlight the current scientific information on CBG, focusing on its pharmacokinetic and pharmacodynamic characteristics and on its principal pharmacological effects, and on pharmacological effects of CBG-derived compounds.

2. Methods

Bibliographic research was carried out independently by two researchers using major scientific databases (PubMed, Scopus, and Google Scholar) using the keyword “cannabigerol” is discussed in the present article. The investigators used the keyword “cannabigerol.” Articles written in the English language and published in peer-reviewed scientific journals describing the pharmacological aspects of cannabigerol published between January 1964 and December 2021 were collected and discussed.

2.1. Pharmacokinetics of Cannabigerol. The pharmacokinetics of CBG has not been still sufficiently characterized in humans. However, there are known methods to quantify CBG presence in the blood [12], such as the liquid chromatography (LC) method, which seems to be more suitable than the gas chromatography (GC) method to measure cannabinoids, because it allows contemporary analysis of acidic and neutral forms. The reason is that acid forms keep their original form during LC analytical conditions, while they turn into neutral form due to the GC temperature conditions [13] (Figure 2).

CBG has been quantified by LC-tandem mass spectrometry in the oral fluid of frequent and occasional cannabis smokers. The maximum concentration (C_{max}), time to C_{max} (T_{max}), and time of last detection (T_{last}) have been calculated, showing that CBG T_{max} occurs at approximately




FIGURE 2: CBG numbering system.

0.17 h after cannabis intake, with a successive quick decrease. It can be detected either in frequent or occasional smokers. In frequent smokers, CBG C_{max} is greater if inhaled than after the oral intake of cannabis; meanwhile, no difference has been reported between smoked and vaporized cannabis. In frequent smokers, CBG T_{last} occurs significantly later after smoking and vaporizing compared to oral administration, thus suggesting that CBG remains longer in the blood circulation after inhalation compared to the oral route (not clear). Generally, CBG cannot be detected beyond 26 hours after any way of administration. In summary, CBG exhibits higher concentrations in the oral fluid of frequent smokers after inhalation than after the oral intake of cannabis [14].

CBG, in the acidic form, is the forerunner of cannabinoids following condensation of geranyl phosphate and olivetolic acid in *C. sativa*. Its identification has been allowed by chemical analysis of urine samples of *Cannabis* consumers. CBG entry in the body occurs with *Cannabis* smoking and its conjugated form is eliminated with urine, as happens for other neutral cannabinoids. Previous chromatographic analysis of hydrolyzed and trimethylsilylated urine samples found a chemical substance in *Cannabis* consumers' urine extracts, with fragment ions at m/z 425, 465, and 479 at a retention time of 14.19 min, which is presumed to be 4''-hydroxy-CBG or 5''-hydroxy-CBG. This was not found in non-hydrolyzed urine samples, suggesting that CBG is also eliminated in the glucuronidated form [15]. The detection rates of CBG in frequent smokers have been shown to be 100% and 90.9% after smoking and vaporization, respectively. Meanwhile, in occasional smokers, the CBG detection rates were 77.8% and 66.7% after smoking and vaporization, respectively. Following detection in frequent smokers ($p = 0.015$), T_{last} was significantly later than in infrequent smokers in a way not dependent on the inhalation method. CBG detection in constant smokers showed considerably greater C_{max} ($p = 0.046$) and later T_{last} ($p = 0.026$) values following smoking in comparison to vaporization, partially due to ineffective CBG evaporation during vaporization. Following cannabis intake, CBG is traceable in the urine where it could be used as an index of current use after inhalation [16].

CBG is considered a potent competitive inhibitor of anandamide (AEA). As occurs for other chemicals, the flexible nature of CBG allows it to occupy most of the active site volume, which could explain its competitiveness versus AEA. CBG competitively inhibits AEA metabolism with a K_i of $10.8 \pm 1.4 \mu M$, whereas THC, $\Delta 8$ -THC, CBD, cannabinol

(CBN), and cannabichromene (CBC) behave as noncompetitive AEA inhibitors. Among these substances, THC is the most powerful inhibitor of the AEA metabolism and causes a reduction of the CYP2J2-mediated AEA metabolic transformation to 20% of the uninhibited activity [17].

CBG, as occurs for $\Delta 9$ -THC, $\Delta 8$ -THC, CBC, CBD, and CBN (Figure 1), is hydroxylated by the isoenzyme CYP2J2. Metabolite formation has been investigated in mice, rats, cats, guinea pigs, hamsters and gerbils, and rabbits. It has been observed that each species is responsible for the formation of similar substances, even if ratios of single metabolites are significantly different [18]. The research identified twelve products of metabolism, among these ones the most important being monohydroxy compounds showing the hydroxyl group at C-8', C-9, C-4', or at positions of the pentyl chain. Metabolites are similar to products of metabolism detectable by investigating other cannabinoids and hydroxylation and epoxidation are the most significant *in vitro* pathways. The principal metabolic pathway in most species is generally allylic hydroxylation of the trans-terminal methyl group (C-8') [19]. A previous study of CBG metabolism in rats, guinea pigs, rabbits, and hamsters showed that this cannabinoid receives allylic hydroxylation at the terminal's double-bond of the C-10 chain, producing the principal metabolites in all species except for mice, where occurs the formation of an epoxide at the terminal's double-bond. In guinea pigs hydroxylation of the pentyl side-chain, particularly at C-3'', was prominent. In other species, 4''-hydroxylation was the undertaken major biotransformation route [20].

The pharmacokinetic profile of CBG after oral and intraperitoneal acute single-dose administration has been investigated in the blood and brains of rats and mice. The maximal drug concentration in tissue (C_{max}), the time to get to the maximal tissue levels (T_{max}), apparent elimination (depending also on the rate of absorption and distribution), half-life (2–6 h after oral administration; per os; 4–24 h after intraperitoneal administration; i.p.), area under the curve (AUC), and brain/plasma ratios founded on the AUC (0–6 h) were measured in the blood and brain specimens at six-time points after i.p. and oral CBG. Rapid absorption was observed both for i.p. and oral CBG administration in the plasma and brain of mice. CBG i.p. injections produced significantly higher plasma and brain concentrations in comparison to oral intake and T_{max} was detected at 30 min in the plasma and 120 min in the brains, while the AUCs were approximately 100-fold greater in comparison to oral way of administration. However, after i.p. injections, the concentrations in both tissues did not show any considerable differences; after oral administration, the concentrations were extremely low and the brain/plasma ratios were 0.77 and 0.15 for the oral and i.p. way of administration, respectively. Moreover, in rats, CBG was quickly absorbed following i.p. and oral administration in both the plasma and the brain, with T_{max} between 30 and 120 min, but i.p. administration provided better exposure, leading to a lower concentration detected following oral intake and stable brain exposure 2–4 h after treatment, with i.p. versus per os brain/plasma ratios overlapping [21].

AEA is converted by different CYP epoxygenases, including CYP2J2, into 14, 15-epoxyeicosatrienoic acid-ethanolamides (EET-EAs) [22]. It has been observed that phytocannabinoids can reduce the metabolism of endogenous substrates such as AEA by inhibiting CYP2J2 [23].

Summarizing the pharmacokinetic research on CBG, it is possible to assert that after inhalation of cannabis, CBG is present in the plasma after a few minutes and reaches T_{max} at approximately 0.17 h, followed by a rapid reduction in concentration. It is absorbed rapidly after i.p. and per os administration. After a single oral administration, its half-life is 2–6 h, and it has also been found in the brain. CBG undergoes a metabolic transformation in the liver, and the major metabolic route is allylic hydroxylation. CBG is metabolized by the enzyme CYP2J2 to produce new hydroxyl and de-oxygenated products. The major metabolites are monohydroxy compounds. CBG is excreted with urine in the conjugated form, as occurs for other cannabinoids.

2.2. Pharmacodynamics of Cannabigerol. Cannabinoid receptors CB1R and CB2R are both associated with inhibitory Gi/o proteins to adenylyl cyclase, and with the mitogen-activating protein (MAP) kinase in an activating way [24]. After receptor activation by phytocannabinoids, the cyclic adenosine monophosphate (cAMP) level is reduced by inhibitory adenylate cyclase and stimulatory MAP kinase. This causes the shortening of the presynaptic action potential, thus inhibiting Ca²⁺ entry and activating the enzymes phospholipases A and C, thus leading to a decrease in excitatory and inhibitory transmitter discharge [25].

CBG binds itself to CB1R and CB2R [26] and is considered a partial agonist at these receptors (K_i values ranging from 300–500 nM) [27]. Other authors have demonstrated that CBG is a poor ligand, binding weakly without activation of CB1R and CB2R, while a sesquicannabigerol, a farnesyl prenylogue of cannabigerol, has been reported to have a higher affinity at CB2R [28]. Other experiments have shown the binding of CBG to CB1R ($K_i = 381$ nM) in mouse brain membranes and CB2R ($K_i = 2.6$ μ M) in CHO cells [29]. It has been shown that CBG binding to CB2R produces a less efficacious CB2R-mediated inhibition of cyclic AMP than THC [27].

CBG shares a similar mechanism of action with the cannabinoid CBD, since it binds itself weakly to CB1R and CB2R and, in addition to blocking CB1R, antagonizes 5-HT_{1A} receptor, activates alpha (2) adrenoceptors [30, 31], and is a modulator of endocannabinoid signaling [30]. CBG acts as a strong α 2-adrenoceptor agonist ($EC_{50} = 0.2$ nM), while it is only a moderate 5-HT_{1A}R competitive antagonist, and at high concentrations (10 μ M), it antagonizes [35S] guanosine 5'-O-[gamma-thio]triphosphate (GTP γ S) binding in mouse brain membranes stimulated with AEA or CP-55940 [29, 30].

It has also been demonstrated that CBG causes inhibition of the reuptake of AEA and acts as an agonist in the TRP channels TRP ankyrin (A) 1, TRP vanilloid (V) 1, and TRPV2, while is an antagonist of the TRP subfamily melastatin (M) 8. Previous findings have shown that CBG can

act as an agonist/desensitizer of transient receptor potential (TRP) ankyrin (A) 1 ($EC_{50} = 700$ nM), an agonist of TRP (vanilloid) V1 ($EC_{50} = 1.3$ μ M), an agonist of TRPV2 ($EC_{50} = 1.7$ μ M), an antagonist of TRP melastatin (M) 8 channels ($IC_{50} = 160$ nM), and an inhibitor of AEA cell uptake ($K_i = 11.3$ μ M) [32]. CBG may also modulate the activity of TRP of ankyrin type-1 channels; however, the EC_{50} values lie in the micromolar range [33]. CBG is an antagonist of TRPM8, useable for possible employment in the treatment of prostate cancer, bladder overactivity, and bladder pain [34, 35].

The peroxisome proliferator-activated receptor (PPAR) γ has also been indicated as a potential pharmacological target of CBG ($K_i = 11.7$ μ M), and concentrations of CBG between 10 and 25 μ M produce enhancement of PPAR γ transcriptional activity [36, 37]. CBG also blocks the voltage-gated sodium channels Nav 1.1, 1.2, and 1.5 and can reduce the psychotropic effect of THC [38].

Both cannabinoids and endocannabinoids possess anti-inflammatory activity through a not sufficiently clarified mechanism of action. It has been suggested that anti-inflammatory activity is a key factor in the development of colon cancer. COX-mediated anti-inflammatory activity of CBG has been studied by using a concentration of $2.5 \cdot 10^{-5}$ M, while CBGA has been tested at a concentration of $6.25 \cdot 10^{-5}$ M. Both CBG and CBGA showed inhibition of the enzyme COX-1 more than a 30%. The same percentage (30%) of inhibition was shown by CBG and CBGA on COX-2 enzyme activity. Anyway, CBG and CBGA inhibition of prostaglandin production resulted to be low being less than 10%. In conclusion, experiments have shown that CBG inhibits COX enzymes in a higher concentration range, compared to conventional anti-inflammatory compounds [39].

3. Biological Effects

3.1. Anti-Inflammatory and Antioxidant Activities. The endocannabinoid system modulates homeostatic processes, including gastrointestinal motility, hunger, pain perception, and immunity [40]. Anti-inflammatory and anti-oxidant activities were investigated mainly through pre-clinical research. In laboratory animals investigation, cannabinoids have been shown to have anti-inflammatory effects, and a subgroup analysis suggested that in experimental colitis, CBG caused the largest reduction in myeloperoxidase activity and in the effect size (SMD -6.20 ; 95% CI, -9.90 to -2.50), followed in producing these effects by the synthetic CB1 agonist arachidonyl-2-chloroethylamide (ACEA) [41].

Inflammatory bowel disease (IBD) includes chronic relapsing inflammatory disorders Crohn's disease (CD) and ulcerative colitis. The effects of CBG were studied in experimental colitis induced in rodents by intracolonic dinitrobenzene sulphonic acid (DNBS). CBG (1–30 mg/kg) i.p. administration before or following induction of experimental colitis significantly improved the consequences of DNBS administration on colon weight/length ratio, an index used for the evaluation of tissue edema caused by inflammation [42]. The effects of CBG on intestinal inflammation

have also been studied by assessing myeloperoxidase activity, histological evaluation and immunohistochemistry, measurement of interleukin (IL)-1 β , IL-10 and interferon (IFN)- γ levels by ELISA, inducible nitric oxide synthase (iNOS), and COX-2, CuZn-superoxide dismutase (SOD) activity. Nitric oxide production and oxidative stress have been evaluated by measuring the effects of CBG on murine macrophages and intestinal epithelial cells, respectively. The results showed that CBG reduces the weight/length ratio of the colon, myeloperoxidase activity, and iNOS expression, and these results were associated with increased SOD activity and normalization of the IL-1 β , IL-10, and IFN- γ modifications caused by DNBS administration. In macrophages, CBG caused reduction of nitric oxide production and iNOS protein (but not mRNA) expression. Interestingly, the CB1R antagonist rimonabant did not modify CBG effects on nitric oxide formation, while the CB2R antagonist SR144528 enhanced the inhibition of CBG on nitric oxide production. According to the authors' conclusions, these experiments show that CBG treatment is able to reduce experimental colitis through a reduction in nitric oxide formation in macrophages via CB2R activation and the production of a diminution of reactive oxygen species (ROS) in intestinal epithelial cells [43].

Other authors have confirmed that cannabinoids inhibit oxidative and nitrosative stress through the modulation of iNOS expression and reduction of ROS [44, 45], and that CBG exerts antioxidant activity comparable to that of vitamin E [46]. The antioxidant activity counteracting hydrogen peroxide (H₂O₂)-induced oxidative stress in murine RAW264.7 macrophages incubated with CBG has been investigated. It was found that 10 μ M of CBG inhibited oxidative stress after stimulation of CB2R, since pre-treatment with the CB2R antagonist AM630 antagonized the protective effects of CBG occurring in H₂O₂-stimulated macrophages, while the involvement of CB1R seems to be excluded. CBG antioxidant activity implies the downregulation of oxidants signals iNOS, nitrotyrosine, and PPAR-1 by preventing I κ B- α phosphorylation and the translocation of nuclear factor- κ B (NF- κ B), as well as the regulation of the MAP kinase pathway. Contextually, it has been found that CBG increases cellular antioxidant defense by modulating SOD-1 expression, thus inhibiting cell death [47]. This indicates that CBG could be useful as a new approach in the care of oxidative stress-related disorders. CBG may effectively work as a free radical scavenger to enhance cellular antioxidant activity through the modulation of pathways such as MAPK kinase and NF- κ B translocation and to counteract cell death.

In the view that inflammation and oxidative stress also have a key role in neurodegeneration, it has been investigated whether CBG may have neuroprotective effects counteracting inflammation and oxidative stress, thus exerting protection against neuronal loss. The capability of CBG to defend neuroblastoma spinal cord (NSC)-34 motor neurons from the toxicity induced in lipopolysaccharide (LPS)-stimulated RAW264.7 macrophages has been evaluated.

Through an *in vitro* MTT [3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyl-2H-tetrazolium bromide] assay (to measure cellular metabolic activity as an indicator of cell viability, proliferation, and cytotoxicity), it was observed that CBG (1–20 μ M) reduced the loss of cell viability induced by LPS-stimulated macrophages in NSC-34 cells. In these experiments, CBG pre-treatment reduced apoptosis, as revealed by a decrease in caspase 3 activation and nuclear-encoded protein Bax expression, while the anti-apoptotic protein Bcl-2 was more elevated. Moreover, CBG pre-treatment reduced inflammation, as demonstrated by the reduction in the IL-1 β , tumor necrosis factor- α (TNF)- α , IFN- γ , and PPAR protein levels evaluated by immunocytochemistry, but also oxidative stress in NSC-34 cells treated with the medium of LPS-stimulated RAW264.7. These experiments suggest that CBG may be a potential treatment to be used against neuroinflammation and oxidative stress [48].

3.2. Antitumoral Activity. Effects of CBG were studied on several tumor cell lines. The antitumoral activity of CBG against human oral epithelioid carcinoma cells has been investigated. In these *in vitro* experiments, CBG showed an antitumoral property against skin melanoma cells with significant activity (IC₅₀ = 31.30 μ M) in an *in vitro* MTT (3-(4, 5-Dimethylthiazol-2-yl)-2, S-diphenyl-2 H-tetrazoliumbromide) microculture assay [49]. The same authors observed that CBG inhibited the growth of mouth epidermal carcinoma cell lines, producing an IC₅₀ of 31 μ M in an MMT assay and 77 μ M in a sulforhodamine B protein assay. In these experiments, CBG showed the least cytotoxic effect on murine NIH 3T3 cells and was more potent than 5-fluorouracil [50]. CBG was found to be effective against breast cancer in the human MDA-MB-231 cell line [51] and to antagonize proliferation of the hyper-proliferating human keratinocyte cell line [52]. Furthermore, as observed above, CBG is able to reduce experimental intestinal inflammation, which might be important in light of the risk of developing colorectal cancer (CRC), which is considerably more elevated in ulcerative colitis patients [53]. From this point of view, the effects of CBG on CRC cell growth and its potential preventive activity in an azoxymethane model of colon cancer, as well as its potential therapeutic effect in a xenograft model of colon cancer, were evaluated. In this experimental model, it was shown that CBG promoted apoptosis, stimulated ROS production, upregulated the mRNA of transcription factor CCAAT-enhancer-binding protein homologous protein (CHOP), and slowed down the cell growth in CRC cells. The effects of CBG on cell growth were not dependent on TRPA1, TRPV1, and TRPV2 channel activation, they were additionally enhanced by the CB2R antagonist SR144528, and were produced also by other TRPM8 channel blockers but not by the 5-HT1A antagonist rimonabant. CBG activity on cellular growth and CHOP mRNA expression was diminished in the TRPM8 cellular line. CBG antagonized the development of xenograft cancers and colon tumors induced by chemicals. In the same experimental model, CBG hampered tumor progression and inhibited the progress of CRC cells [54]. These *in vivo*

experiments show that CBG antagonizes the growth of CRC cells principally by stimulating apoptosis and hindering the development of colon carcinogenesis. Moreover, authors have suggested that TRPM8 is also at least partially involved in the antitumoral activity of CBG. The suppressor activity of CBG of the growth on tumor cell lines is accompanied by ROS hyperproduction and seems to be partially mediated by TRPM8. The apparent safety and experimental results of CBG indicate that it could be promising for preventing and treating CRC.

Glioblastoma is aggressive brain cancer, showing increasing incidence. It has been found that CBG, as occurs with THC, slows the progression of this tumor and inhibits the invasion of glioblastoma cells. Furthermore, CBG is effective in reducing therapy-resistant glioblastoma stem cells [55]. These findings indicate that CBG, also showing analgesic, anti-nausea, and appetite-stimulating effects, could represent a new adjuvant therapy for glioblastoma.

3.3. Neuroprotective Effects. CBG as a cannabis-derived chemical not causing psychotomimetic effects is interesting as a potential new drug for central nervous system (CNS) pathologies.

Neuroprotective effects of CBG and CBD have been compared in experiments simulating oxidative stress and neurotoxicity as they occur in neurological pathologies in rats. CBG and CBD exert antioxidant activity in astrocytes exposed to hydrogen peroxide and restored the content of serotonin in the cortex [56].

The effects of CBG have also been studied on human brain microvascular endothelial cells (HBMECs), pericytes, and astrocytes forming the blood-brain barrier (BBB) under ischemic conditions. Through this experimental model, it has been shown that 10 μ M of CBG reduced IL-6, lactate dehydrogenase, and DNA damage protein levels in astrocytes [57].

The protective effects of CBG have been investigated in two *in vivo* models of Huntington's disease (HD) induced by 3-nitropropionate (3NP). CBG administered through daily four i.p. injections (10 mg/kg) in mice improved motor deficits, preserved striatal neurons, attenuated microgliosis, and reduced inflammatory markers induced by 3NP against 3NP toxicity, and ameliorated antioxidant activity. CBG also produced significant recovery in the deteriorated rotarod performance. Furthermore, by using HD array analysis, it was evident that CBG turned to normal a series of genes related to this pathology [58].

The synthetic analog cannabigerol quinone VCE-003 (Figure 3), obtained by oxidation modification in the resorcinol moiety of the compound, has been identified as an anti-inflammatory substance. VCE-003 reduced the effects of excitotoxicity on neuronal cells, activated PPAR γ transcriptional activity, and antagonized the release of pro-inflammatory substances in an experimental model characterized by the stimulation of microglial cells with lipopolysaccharide (LPS). In the same series of experiments, the authors found that VCE-003 improved the symptoms of

a model of multiple sclerosis (MS) represented by Theiler's murine encephalomyelitis virus (TMEV) infection [59].

Multiple sclerosis (MS) is a chronic inflammatory demyelinating CNS disease, considered a predominantly T cell-mediated autoimmune disease and neuroinflammation is a fundamental component in the pathophysiology of this disease [60]. It has been demonstrated that VCE-003, reduces neuroinflammation and motor deficiency in viral T cells and macrophages, and its effectiveness in an autoimmune model of MS has been investigated. Proliferation, cell cycle, and expression of activation markers were assessed by fluorescence-activated cell sorting analysis in human primary T cells, and cytokine and chemokine formation was quantified. In the same study, transcription was investigated by using Jurkat and RAW264.7 cells with the aim of evaluating VCE-003 activity on IL-17-induced macrophage polarization to activate the M1 phenotype. Experimental autoimmune encephalomyelitis (EAE) was induced by myelin oligodendrocyte glycoprotein (MOG35–55) immunization and spinal cord harm was established with immunohistochemistry. The results showed that the postimmunization administration of i.p. VCE-003 (5 mg/kg) for 21 consecutive days inhibited proliferative effects stimulated by CD3/CD28, cell cycle progression, and induction of IL-2Ra and ICAM-1 in human T cells. VCE-003 also inhibited the formation of Th1/Th17 cytokines and chemokines in murine T cells. This effect was associated with a reduction of transcription of IL-2, IL-17, and TNF- α promoters when induced by CD3/CD28. Moreover, VCE-003 and the CB2 agonist JWH-133, attenuated the pro-inflammatory polarization of macrophages caused by IL-17. Another effect of VCE-003 was the prevention of iNOS expression stimulated by LPS. VCE-003 ameliorated the neurological profile and harshness of EAE in mice by stimulation of CB2 and PPARc receptors. Finally, in these experiments, attenuation of cell infiltrates, associated with reduction of microglial activation, structural conservation of myelin, and protection of axons, were observed. Thus, indicating the potential role of VCE-003 as a therapeutic compound to use in immune diseases characterized by inflammation [61].

The analog VCE-003.2 (Figure 3), another CBG quinone derivative, acting through PPAR γ (a nuclear receptor involved in the lipidic metabolism and glucose homeostasis), has been studied for its potential neuroprotection. It has been shown that it favors the pro-survival of progenitor cells all along with neuronal differentiation, and this effect was interfered with by PPAR γ antagonists. Moreover, VCE-003.2 given i.p. at the dose of 10–20 mg/kg reduced cell death caused by quinolinic acid (QA), activation of caspase-3 activation, and accumulated mutant huntingtin aggregates in cells originating from the striatum. In the *in vivo* Huntington's-like disease models characterized by striatal neurodegeneration induced by QA and 3NP, VCE-003.2 prevented medium spiny DARPP32+ neuronal loss, ameliorated motor deficits, reactive astrogliosis, and microglial activation. In the 3NP model, VCE-003.2 antagonized the formation of proinflammatory markers and improved brain antioxidant activity. Taken together, the results of these




FIGURE 3: CBG pleiotropic mechanism of action.

experiments bring to consider the compound VCE-003.2 as a potential candidate for the therapy of HD and other neurodegenerative pathologies characterized by the presence of inflammatory components [62].

Neuroprotection of VCE-003.2 in amyotrophic lateral sclerosis (ALS) was investigated by using SOD1G93A mutant mice. VCE-003.2 (10 mg/kg) was administered i.p. from 60 days up to 18 weeks when the disease is in an advanced stage. Treatment with VCE-003.2 decreased neurological abnormalities, protected spinal cholinergic motor neurons, and attenuated astrogliosis. Elevation in IL-1 β and glial glutamate transporters, as well as the LPS-induced release of TNF- α and IL-1 β in cultured astrocytes of SOD1G93A transgenic newborns, were reduced, probably through PPAR- γ stimulation. The results showed the neuroprotective effects of VCE-003.2 in SOD1G93A transgenic mice, indicating PPAR- γ as an additional therapeutic target suitable for the utilization of cannabinoids in ALS [63].

The effects of VCE-003.2 were investigated *in vivo* in mice in which unilateral intrastratial lesions were caused by the injection of LPS to reproduce a model of Parkinson's disease (PD), as well as in an *in vitro* model (LPS-exposed microglial BV2 cells and immortalized rat embryonic striatal M-213 cells treated with media obtained from LPS-exposed BV2 cells). I.p. administration of VCE-003.2 (10 mg/kg), alone or in association with the PPAR γ antagonist T0070907 (5 mg/kg) given for 21 days beginning 16 h after LPS lesion, decreased the loss of nigrostriatal neurons and the intense microgliosis in the *Substantia nigra*, measured by Iba-1/Cd68 immunostaining. Mediators of inflammation TNF- α , IL-1 β , and iNOS in the striatum showed marked elevation by the LPS lesion and were significantly reduced by VCE-003.2 via activation of PPAR γ . Moreover, the *in vitro* approach confirmed the anti-inflammatory activity of VCE-003.2, showing that in LPS-exposed BV2 cells it reduced the synthesis and release of TNF- α and IL-1 β , and the induction of iNOS and COX-2 [64].

More recently, VCE-003.2 has been investigated in mice in another model of PD, in this case, induced with the neurotoxin 6-hydroxydopamine (6-OHDA), comparing its effects with two other CBG-related derivatives,

cannabigerolic acid quinone (CBGA-Q) and its sodium salt (CBGA-Q-Salt). These compounds, as well as VCE-003.2, act at PPAR- γ receptor, but are not ligands for CB1R and CB2R. VCE-003.2 was cytoprotective in the SH-SY5Y cell line exposed to 6-OHDA at a concentration of 20 μ M, an effect not changed by the PPAR- γ receptor antagonist GW9662. Cytoprotection was also observed with CBGA-Q and CBGA-Q-Salt; however, this effect was abolished by GW9662. *In vivo* experiments showed that the oral administration of VCE-003.2 (20 mg/kg) protected nigral neurons against 6-OHDA-induced damage. This neuroprotection was accompanied by motor deficiencies caused by 6-OHDA. Similar protection was observed with CBGA-Q, given orally (20 mg/kg) or intraperitoneally (10 mg/kg, i.p.), but to a lower extent, while CBGA-Q-Salt (10 mg/kg) was weakly active. The authors' interpretation emphasized the role of PPAR- γ receptors involvement in these effects [65].

The protective effects of VCE-003.2 have also been studied in striatal neurodegeneration by using *in vivo* adeno-associated viral expression of mutant huntingtin and *in vitro* mouse embryonic stem cell differentiation. The effects of VCE-003.2 on embryonic stem-neuronal differentiation were investigated. The R1 line of mouse ES cells was treated with VCE-003.2 during neural differentiation for 21 days and transcription factor CTIP2-positive striatal medium spiny neurons were examined. Immunofluorescence showed that VCE-003.2 enhanced the quantitative of CTIP2-positive cells as well as doublecortin immunoreactivity and favored differentiation. In *in vivo* experiments, daily orally administered VCE-003.2 (10 mg/kg) protected striatal neurons from injury caused by mutant huntingtin, reduced neuroinflammation, and ameliorated motor deficit. Other effects of VCE-003.2 were the promotion of mobilization of sub-ventricular zone progenitor and of migration of neuroblasts toward the damaged area, and increase in neurogenesis. VCE-003.2 also increased neuroblasts and striatal-like neurogenesis. In summary, VCE-003.2 improved sub-ventricular zone-derived neurogenesis against neurodegeneration caused by mutant huntingtin, and was shown to be neuroprotective upon oral administration [66].

3.4. Anti-Anxiety Effects. Endocannabinoids have been shown to modulate feeding and emotional behaviors [67, 68]. CBG acts as a 5-HT_{1A} antagonist as demonstrated in *in vitro* [30] and *in vivo* experiments [69]. For this reason, it has been proposed that it can be responsible for the anti-anxiety effects of cannabis [70]. However, the notion that CBG behaves as a 5-HT_{1A} antagonist conflicts its putative anxiolytic effects because an agonist—or partial agonist—would be more likely to produce anxiolytic effects. These aspects were investigated by comparing the effects of acute and chronic (14 days) i.p. CBG (2.5 mg/kg) with the effects of chronic i.p. administration of THC (2.5 mg/kg) and CBD (2.5 mg/kg) on the time consumed in an open, lit box in a light–dark (LD) immersion model of anxiety-like behavior and saccharin hedonic reactions in a taste reactivity test of palatability processing in rats. The results showed that THC caused acutely an anxiogenic-like behavior in the LD immersion test, not enhanced by chronic administration. THC increased only transiently on the first day of administration the saccharin palatability in the taste reactivity test. Both CBD and CBG did not change the anxiety-like response but CBG produced a light increase of saccharin palatability only on the first day of administration. These results were unable to corroborate the anti-anxiety effects of CBG in rats because they did not show the anxiolytic effect of CBG [71]. Contrary to these results, in other experiments, it has been observed that i.p. CBG administration at a dose of 10 mg/kg in mice enhanced the period of time consumed in the central quadrant of the open field test, thus suggesting potential anxiolytic effects. In the same group of experiments, i.p. CBG administration at a dose of 3 mg/kg also produced a light anti-nociceptive effect [72].

3.5. Anti-Nausea Effects. CBG seems to influence the anti-emetic properties of THC. Antagonism of CBG toward CB₁R and 5-HT_{1A} receptors [30] abolishes the anti-emetic activity of low-dose of CBD, probably because it is due to action at the 5-HT_{1A} receptor level [64]. The potential of CBG to antagonize the anti-emetic effect of the 8-OH-DPAT antagonist of the 5-HT_{1A} receptor [73], has also been assessed. The emetic effects properties of higher doses of CBD and CBG may have involved in the production of severe nausea and vomiting when cannabinoid hyperemesis syndrome occurs [74]. Experiments have been conducted to study and evaluate possible CBG effects either in regulating nausea in rats or vomiting in *Suncus murinus*. Two experiments were carried out; in experiment 1, rats received i.p. CBG (0.0, 1, 5, and 10 mg/kg) 15 min prior to receive i.p. vehicle or CBD (5 mg/kg; experiment 1a) or i.p. vehicle or 8-OH-DPAT (0.01 mg/kg; experiment 1b). After 30 min, all rats were i.p. treated with the association of 0.1% saccharin solution and LiCl (20 mL/kg of 0.15 M). Seventy-two hours later, animals were subjected to a saccharin taste reactivity to observe the effects on the establishment of conditioned gaping reactions (an experimental model of nausea) and conditioned saccharin avoidance. In experiment 2, *S. murinus* were i.p. treated with CBG (5 mg/kg) or vehicle 15 min before CBD (5 mg/kg) or vehicle, and 30 min later i.p.

treated with LiCl (60 mL/kg of 0.15 M), to evaluate the number of vomiting episodes. CBD blocked conditioned gaping in rats and vomiting in shrews, and these effects were antagonized by pretreatment with all doses of CBG. CBG also antagonized the anti-nausea effects of 8-OH-DPAT. These results indicate that injection of moderate doses of CBG and CBD may be opposite to each other at the 5-HT_{1A} receptor level with interference in the modulation of nausea and vomiting. These findings suggest that CBG could be used for its potential interactions with CBD due to its opposite action at the 5-HT_{1A} receptor in nausea and vomiting [69].

3.6. Reduction in Intraocular Pressure (IOP). A reduction of IOP in humans following smoked cannabis was noted in 1971 [75]. Successively, findings carried out in different species have shown that THC has an ocular hypotensive effect [76]. Cannabinol (CBN) or CBG have been studied in cats by topically administering them doses of 250, 500, and 1000 picograms as a single drop or through via osmotic minipumps (20 µg/h) for nine days. CBN single dose had a moderate effect on IOP, but it produced a significant reduction if chronically administered. Similarly, CBG reduced IOP, but the size of the effect when it was chronically administered was greater. Furthermore, CBN but not CBG caused side effects such as conjunctival erythema and hyperemia. Following i.p. administration of CBN (20, 40, or 80 mg/kg) to rats, 8–13 Hz polyspike discharges were observed in the electrocorticogram during vigilance and sleep rapid eye movement episodes. Interestingly, systemically injected CBG (10, 30, and 100 mg/kg) lacked this effect. The above-cited results suggest that chronic administration of these cannabinoids reduces ocular tension. However, as happens for cannabis and THC, CBN can cause both ocular toxicity and neurotoxicity. As CBG did not produce these toxicities, it is evident that the ocular hypotensive effect obtained with this cannabinoid can be distinguished from both the adverse central and ocular effects associated with cannabis [77]. When CBG was chronically unilaterally delivered for 9 consecutive days to the eyes of cats through osmotic minipumps releasing it at a rate of 1 µL/h, it produced a fall in IOP similarly to THC (4–7 mm Hg) applied in the same way. The same group of researchers observed that in rats, following peripheral administration of THC, but not CBG, polyspike discharge became visible in the cortical electroencephalogram during the wake. Then, polyspikes were detectable within rapid eye movement sleep periods. Results also showed that both cannabinoids caused a two- to three-fold increase in aqueous outflow [78]. As the greatest proportion of the drugs used for the therapy of glaucoma suppresses aqueous release, CBG could be a promising anti-glaucoma substance to be used with drugs acting by enhancing aqueous outflow instead by increasing of ocular fluid drainage.

IOP-lowering activity was studied in rats using a synthetic CBG analog, cannabigerol-dimethyl heptyl (CBG-DMH), compared to the nonselective CB₁R and CB₂R agonist, WIN55, 212-2. CBG-DMH i.p. administration reduced IOP at doses of 2.5 mg/kg or given locally (1%–2%).

CBG-DMH reducing IOP were antagonized by i.p. administration of O-1918 (2.5 mg/kg), a selective antagonist of the abnormal CBD-sensitive cannabinoid-related receptor (CBx), but not influenced by the CB1R antagonist AM251 (2.5 mg/kg) or the CB2R antagonist AM630 (2.5 mg/kg). Contemporary administration of WIN55, 212-2 given at a subthreshold dose to lower IOP (0.25 mg/kg), in association with topical CBG-DMH (0.25%), augmented the IOP-lowering effect of the compound applied alone. These data show that an analog of CBG decreases IOP in normotensive rat eyes independently of CB1R or CB2R involvement, probably activating putative cannabinoid receptors. The greater reduction in IOP seen with co-application of the CB1R agonist WIN55, 212-2 and CBG-DMH, further indicates that effects on the eye induced by CBG-DMH are involving biological targets different from CB1R [79]. Combined, these findings show that CBG may reduce IOP, with this latter study suggesting that synthetic CBG can decrease IOP through mechanisms not involving CB1R or CB2R activation. This finding implies the existence of putative cannabinoid receptors regulating aqueous humor outflow and probably localized to the anterior ocular tissues.

3.7. Effects on Skin. The effects of CBG on the expression of keratins 1 and 10, involucrin, and transglutaminase 5, and on DNA methylation of keratin 10 gene, have been studied in the human keratinocytes (HaCaT) cellular line, together with DNA methylation and expression of four DNA methyltransferases (DNMT1, 3a, 3b, and 3L). These experiments showed that CBG caused a significant reduction in the expression of the genes investigated by increasing the DNA methylation of the gene for keratin 10. The data obtained from these experiments led the authors to believe that CBG behaves as a transcriptional suppressor, able to control cell proliferation and cell differentiation, being a substance potentially useful for new therapeutic approaches for skin disorders [80]. On this basis, CBG has been studied as a possible anti-acne drug. Acne is a common skin pathology; however, its more serious expression can deeply weaken the quality of life and, because of social censure, can produce consequent psychological disturbance [81]. The anti-acne effects of CBG on the viability and proliferation of sebocytes were observed through MTT and CyQUAN methods. CBG in a significant way reduced cell viability after a 24 h treatment, but not with treatment lasting 48 h. CBG (10–20 mM) treatment for 24 h significantly reduced lipogenesis stimulated by AEA. These results suggest the possibility that CBG may act as a partial agonist through the same pro-lipogenic signaling pathway on which AEA is active. CBG suppressed inflammation caused by LPS on sebocytes. The results of the above-exposed experiments according to the antiproliferative action of phytocannabinoids, increase the hypothesis that these compounds may be useful in acne and in other skin pathologies associated with inflammatory characteristics, such as psoriasis [82].

3.8. Other Effects. It has been suggested that some *C. sativa* derived products might exert positive influence on lower

urinary tract symptoms. Similarly, to THC and CBD, CBG is also able to reduce bladder contractility. Experiments with CBG have been conducted on mouse and human isolated bladders. In particular, CBG at concentrations ranging from 10^{-8} to 10^{-4} M, reduced the contractions of mouse bladders induced by acetylcholine, without influencing bladder contractions induced by electrical field stimulation. The results showed that neither rimonabant (10^{-6} M) nor SR144528 (10^{-7} M), selective CB1R and CB2R antagonists, modified the inhibitory effect of CBG. CBG also inhibited acetylcholine-induced contractions of human bladders, the effect being significant for the 3×10^{-5} M and 10^{-4} M concentrations. These findings indicate that CBG is a more promising compound with respect to other cannabinoids and that it inhibits bladder contraction through a post-synaptic action. The possible involvement of cannabinoid receptors has been hypothesized on the basis of the CBG partial agonism toward CB1R and CB2R and the known effect of CBG on the inhibition of the reuptake of the endocannabinoid anandamide, and finally because components of the endocannabinoid system have a role in the regulation of bladder function. However, the involvement of cannabinoid receptors has not been demonstrated [83, 84].

CBG inhibits platelet aggregation and [14 C]5-HT release. In a series of experiments, CBG did not inhibit tetradecanoylphorbol acetate (TPA)-induced aggregation of human or rabbit platelets; however, in a dose-dependent way (10^{-3} to 10^{-5} M), it partially inhibited primary aggregation (due to the direct interaction of the aggregating agent with its receptor) and totally inhibited secondary aggregation induced by adrenaline. CBG, dose-dependently also inhibited rabbit and human platelet aggregation stimulated by adenosine diphosphate (IC₅₀ of CBG being $2 \times 7 \times 10^{-4}$ M) and in rabbit platelets by platelet-activating factor (concentration range to 3×10^{-5} M and 10^{-4} M) [85].

CBG and other major cannabinoids such as CBD, CBC, THC, and CBN showed activity *versus* methicillin-resistant *Staphylococcus aureus* (MRSA) strains. The rare occurrence of cross-resistance between microorganisms and plant antibacterial-derived compounds suggests the possibility that CBG could be studied as a potential source of compounds to address antibiotic resistance [86]. CBG also showed activity against *Streptococcus mutans* by altering bacterial membrane properties through the induction hyperpolarization and by reducing membrane fluidity and causing a consequent increase in membrane permeability [87]. Other experiments have demonstrated that CBG is a potential agent against bacterial biofilm, a key factor for contamination of medical devices, favoring the raising of human chronic infections [88].

The appetite-stimulating effects of *C. sativa* have been prevalently assigned to THC, while CBG is also an appetite stimulator. CBG given per os in rats at a dose of 30–240 mg/kg induces an increase in appetite-feeding behavior without any influence on motor activity [89, 90]. Since the loss of muscle, anorexia, and metabolic dysfunction are common consequences of chemotherapy with cytotoxic drugs, *C. sativa* derivatives were used to reduce these effects in oncologic patients. With this aim, the most used were




FIGURE 4: CBG synthetic analogues.

Cannabis extracts containing significant amounts of THC or synthetic THC analogs. Unfortunately, the effect of THC psychotomimetics represents an obstacle to its utilization. For this reason, on the basis of the appetite-stimulating effect of CBG, it has been assessed in laboratory animals whether this cannabinoid given orally for three days reduces anorexia and/or other cachectic signs induced by the i.p. administration of the chemotherapeutic substance cisplatin (6 mg/kg). CBG was able to reduce the anorexia, weight loss, and metabolic dysfunction caused by cisplatin [91].

Recently, three new CBG derivatives (HUM-223, HUM-233, and HUM-234) (Figure 4) were synthesized, showing anti-inflammatory and analgesic properties. Additionally, one of them, HUM-234, arrested the development of obesity in mice fed with a high-fat food regimen [92].

4. Conclusion

CBG, in its acidic form, is the precursor of the most well-known cannabis derivative compounds, namely, THC and CBD. The study of its pharmacology shows that this

compound shares some characteristics with other phytocannabinoids, but it displays its own characteristic profile, as shown by emerging research. Pharmacodynamic research suggests a mechanism of action involving only partial activity of classical cannabinoid receptors, indicating that CBG, as well as CBD, has a multitarget pharmacological action and interacts with a number of endocannabinoid and non-endocannabinoid targets (Figure 3). Research, both *in vivo* and *in vitro*, shows several pharmacological effects of CBG, such as a reduction in IOP and dermatological, anti-inflammatory, antioxidant, antitumoral, and anti-anxiety activities, together with appetite-stimulating effects. Research on CBG also shows the development in the production of synthetic derivatives. Synthetic analogs have been tested with positive effects, particularly in the field of neuroprotection. The promising results obtained with CBG, together with the apparent lack of psychotomimetic THC-like effects, suggest that research on CBG deserves to be deepened as it could be used, alone or in association, for novel therapeutic approaches for several disorders.

However, these effects have been proven only through preclinical experiments, and clinical research is needed to confirm these potential activities in humans.

Abbreviation

ALS:	Amyotrophic lateral sclerosis
AEA:	Anandamide
ACEA:	Arachidonyl-2-chloroethylamide
AUC:	Area under the curve
BBB:	Blood-brain barrier
CHOP:	CCAAT-enhancer-binding protein homologous protein
CBG:	Cannabigerol
CBG-	Cannabigerol-dimethyl heptyl
DMH:	
C. sativa:	Cannabis sativa
CBGA:	Cannabigerolic acid
CBGA-Q:	Cannabigerolic acid quinone
CBD:	Cannabidiol
CBC:	Cannabichromene
CBN:	Cannabinol
$\Delta 8$ -THC:	$\Delta 8$ -tetrahydrocannabinol
$\Delta 9$ -THC:	$\Delta 9$ -tetrahydrocannabinol
CYP:	Cytochrome P
cAMP:	Cyclic adenosine monophosphate
CB1R:	Cannabinoid receptor 1
CB2R:	Cannabinoid receptor 2
CNS:	Central nervous system
CRC:	Colorectal cancer
CD:	Crohn's disease
COX-1:	Cyclooxygenase 1
COX-2:	Cyclooxygenase 2
MTT:	[3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyl-2H-tetrazolium bromide]
DNBS:	Dinitrobenzene sulphonic acid
DNMTs:	DNA methyltransferases
EAE:	Experimental autoimmune encephalomyelitis
GTP γ S:	Guanosine 5'-O-[gamma-thio]triphosphate
HBMECs:	Human brain microvascular endothelial cells
HD:	Huntington's disease
H ₂ O ₂ :	Hydrogen peroxide
6-OHDA:	6-hydroxydopamine
5-HT1A:	5-hydroxytryptamine 1A
iNOS:	Inducible nitric oxide synthase
IBD:	Inflammatory bowel disease
IFN- γ :	Interferon
IL:	Interleukin
i.p.:	Intraperitoneal
IOP:	Intraocular pressure
LD:	Light-dark
LPS:	Lipopolysaccharide
LC:	Liquid chromatography
GC:	Gas chromatography
C _{max} :	Maximum concentration
MAP kinase:	Mitogen-activating protein kinase
MS:	Multiple sclerosis
MOG:	Myelin oligodendrocyte glycoprotein

NSC:	Neuroblastoma spinal cord
3NP:	3-nitropropionate
I κ B- α :	NF-kappa-B inhibitor alpha
NF- κ B:	Nuclear factor- κ B
PD:	Parkinson's disease
PPAR γ :	Peroxisome proliferator-activated receptor
QA:	Quinolinic acid
ROS:	Reactive oxygen species
SOD:	Superoxide dismutase
TMEV:	Theiler's murine encephalomyelitis virus
T _{max} :	Time to C _{max}
Tlast:	Time of last detection
TNF- α :	Tumor necrosis factor- α
TRP channel:	Transient receptor potential channels
TRPA1:	TRP ankyrin (A) 1
TRPM8:	TRP subfamily melastatin (M) 8
TRPV1:	TRP vanilloid (V) 1
TRPV2:	TRP vanilloid (V) 2.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

References

- [1] Y. Gaoni and R. Mechoulam, "Structure synthesis of cannabigerol new hashish constituent," *Proceedings of the Chemical Society, London*, vol. 82, pp. 2189–2192, 1964.
- [2] R. Mechoulam and Y. Gaoni, "Hashish—IV," *Tetrahedron*, vol. 21, no. 5, pp. 1223–1229, 1965.
- [3] M. A. Elsohly and D. Slade, "Chemical constituents of marijuana: the complex mixture of natural cannabinoids," *Life Sciences*, vol. 78, no. 5, pp. 539–548, 2005.
- [4] Y. Grunfeld and H. Edery, "Psychopharmacological activity of the active constituents of hashish and some related cannabinoids," *Psychopharmacologia*, vol. 14, no. 3, pp. 200–210, 1969.
- [5] R. Mechoulam, A. Shani, H. Edery, and Y. Grunfeld, "Chemical basis of hashish activity," *Science*, vol. 169, no. 3945, pp. 611–612, 1970.
- [6] E. P. M. de Meijer, M. Bagatta, A. Carboni et al., "The inheritance of chemical phenotype in *Cannabis sativa L.*" *Genetics*, vol. 163, no. 1, pp. 335–346, 2003.
- [7] F. Pollastro, L. De Petrocellis, A. Schiano-Moriello et al., "Amorfrutin-type phytocannabinoids from *Helichrysum umbraculigerum*," *Fitoterapia*, vol. 123, pp. 13–17, 2017.
- [8] C. Citti, B. Pacchetti, M. A. Vandelli, F. Forni, and G. Cannazza, "Analysis of cannabinoids in commercial hemp seed oil and decarboxylation kinetics studies of cannabidiolic acid (CBDA)," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 149, pp. 532–540, 2018.
- [9] E. P. M. de Meijer and K. M. Hammond, "The inheritance of chemical phenotype in *Cannabis sativa L.* (II): cannabigerol predominant plants," *Euphytica*, vol. 145, no. 1-2, pp. 189–198, 2005.
- [10] E. P. M. de Meijer, K. M. Hammond, and A. Sutton, "The inheritance of chemical phenotype in *Cannabis sativa L.* (IV): cannabinoid-free plants," *Euphytica*, vol. 168, no. 1, pp. 95–112, 2009.
- [11] M. Wang, Y. H. Wang, B. Avula et al., "Decarboxylation study of acidic cannabinoids: a novel approach using ultra-high-

- performance supercritical fluid chromatography/photodiode array-mass spectrometry,” *Cannabis and Cannabinoid Research*, vol. 1, no. 1, pp. 262–271, 2016.
- [12] K. B. Scheidweiler, M. N. Newmeyer, A. J. Barnes, and M. A. Huestis, “Quantification of cannabinoids and their free and glucuronide metabolites in whole blood by disposable pipette extraction and liquid chromatography-tandem mass spectrometry,” *Journal of Chromatography A*, vol. 1453, pp. 34–42, 2016.
- [13] B. Patel, D. Wene, and Z. T. Fan, “Qualitative and quantitative measurement of cannabinoids in cannabis using modified HPLC/DAD method,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 146, pp. 15–23, 2017.
- [14] M. J. Swortwood, M. N. Newmeyer, M. Andersson, O. A. Abulseoud, K. B. Scheidweiler, and M. A. Huestis, “Cannabinoid disposition in oral fluid after controlled smoked, vaporized, and oral cannabis administration,” *Drug Testing and Analysis*, vol. 9, no. 6, pp. 905–915, 2017.
- [15] E. Hidvégi and G. P. Somogyi, “Detection of cannabigerol and its presumptive metabolite in human urine after Cannabis consumption,” *Pharmazie*, vol. 65, no. 6, pp. 408–411, 2010.
- [16] M. N. Newmeyer, M. J. Swortwood, A. J. Barnes, O. A. Abulseoud, K. B. Scheidweiler, and M. A. Huestis, “Free and glucuronide whole blood cannabinoids’ pharmacokinetics after controlled smoked, vaporized, and oral cannabis administration in frequent and occasional cannabis users: identification of recent cannabis intake,” *Clinical Chemistry*, vol. 62, no. 12, pp. 1579–1592, 2016.
- [17] W. R. Arnold, A. T. Weigle, and A. Das, “Cross-talk of cannabinoid and endocannabinoid metabolism is mediated via human cardiac CYP2J2,” *Journal of Inorganic Biochemistry*, vol. 184, pp. 88–99, 2018.
- [18] L. O. Hanuš, S. M. Meyer, E. Muñoz, O. Tagliatalata-Scafati, and G. Appendino, “Phytocannabinoids: a unified critical inventory,” *Natural Product Reports*, vol. 33, no. 12, pp. 1357–1392, 2016.
- [19] D. J. Harvey and N. K. Brown, “In vitro metabolism of cannabigerol in several mammalian species,” *Biological Mass Spectrometry*, vol. 19, no. 9, pp. 545–553, 1990.
- [20] D. J. Harvey and N. K. Brown, “Comparative in vitro metabolism of the cannabinoids,” *Pharmacology Biochemistry and Behavior*, vol. 40, no. 3, pp. 533–540, 1991.
- [21] S. Deiana, A. Watanabe, Y. Yamasaki et al., “Plasma and brain pharmacokinetic profile of cannabidiol (CBD), cannabidivarin (CBDV), Δ^9 -tetrahydrocannabinol (THCV) and cannabigerol (CBG) in rats and mice following oral and intraperitoneal administration and CBD action on obsessive-compulsive behaviour,” *Psychopharmacology (Berl)*, vol. 219, no. 3, pp. 859–873, 2012.
- [22] D. R. McDougle, J. E. Watson, A. A. Abdeen et al., “Anti-inflammatory ω -3 endocannabinoid epoxides,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 114, no. 30, pp. E6034–E6043, 2017.
- [23] A. Das, A. T. Weigle, W. R. Arnold, J. S. Kim, L. N. Carnevale, and H. C. Huff, “CYP2J2 molecular recognition: a new Axis for therapeutic design,” *Pharmacology & Therapeutics*, vol. 215, Article ID 107601, 2020.
- [24] A. C. Howlett, F. Barth, T. I. Bonner et al., “International union of pharmacology. XXVII. Classification of cannabinoid receptors,” *Pharmacological Reviews*, vol. 54, no. 2, pp. 161–202, 2002.
- [25] R. G. Pertwee, “The pharmacology of cannabinoid receptors and their ligands: an overview,” *International Journal of Obesity*, vol. 30, no. S1, pp. S13–S18, 2006.
- [26] R. Nachnani, W. M. Raup-Konsavage, and K. E. Vrana, “The pharmacological case for cannabigerol,” *Journal of Pharmacology and Experimental Therapeutics*, vol. 376, no. 2, pp. 204–212, 2021.
- [27] G. Navarro, K. Varani, I. Reyes-Resina et al., “Cannabigerol action at cannabinoid CB (1) and CB (2) receptors and at CB (1)-CB (2) heteroreceptor complexes,” *Frontiers in Pharmacology*, vol. 9, p. 632, 2018.
- [28] F. Pollastro, O. Tagliatalata-Scafati, M. Allarà et al., “Bioactive prenylogous cannabinoid from fiber hemp (*Cannabis sativa*),” *Journal of Natural Products*, vol. 74, no. 9, pp. 2019–2022, 2011.
- [29] R. G. Pertwee, “The diverse CB₁ and CB₂ receptor pharmacology of three plant cannabinoids: Δ^9 -tetrahydrocannabinol, cannabidiol and Δ^9 -tetrahydrocannabivarin,” *British Journal of Pharmacology*, vol. 153, no. 2, pp. 199–215, 2008.
- [30] M. G. Cascio, L. A. Gauson, L. A. Stevenson, R. A. Ross, and R. G. Pertwee, “Evidence that the plant cannabinoid cannabigerol is a highly potent α 2-adrenoceptor agonist and moderately potent 5HT_{1A} receptor antagonist,” *British Journal of Pharmacology*, vol. 159, no. 1, pp. 129–141, 2010.
- [31] R. G. Pertwee, “Receptors and channels targeted by synthetic cannabinoid receptor agonists and antagonists,” *Current Medicinal Chemistry*, vol. 17, no. 14, pp. 1360–1381, 2010.
- [32] L. De Petrocellis, A. Ligresti, A. S. Moriello et al., “Effects of cannabinoids and cannabinoid-enriched Cannabis extracts on TRP channels and endocannabinoid metabolic enzymes,” *British Journal of Pharmacology*, vol. 163, no. 7, pp. 1479–1494, 2011.
- [33] L. De Petrocellis, V. Vellani, A. Schiano-Moriello et al., “Plant-derived cannabinoids modulate the activity of transient receptor potential channels of ankyrin type-1 and melastatin type-8,” *Journal of Pharmacology and Experimental Therapeutics*, vol. 325, no. 3, pp. 1007–1015, 2008.
- [34] L. De Petrocellis and V. Di Marzo, “Non-CB₁, non-CB₂ receptors for endocannabinoids, plant cannabinoids, and synthetic cannabimimetics: focus on G-protein-coupled receptors and transient receptor potential channels,” *Journal of Neuroimmune Pharmacology*, vol. 5, no. 1, pp. 103–121, 2010.
- [35] G. Mukerji, J. Waters, I. P. Chessell, C. Bountra, S. K. Agarwal, and P. Anand, “Pain during ice water test distinguishes clinical bladder hypersensitivity from overactivity disorders,” *BMC Urology*, vol. 6, no. 1, p. 31, 2006.
- [36] X. Nadal, C. Del Río, S. Casano et al., “Tetrahydrocannabinolic acid is a potent PPAR γ agonist with neuroprotective activity,” *British Journal of Pharmacology*, vol. 174, no. 23, pp. 4263–4276, 2017.
- [37] S. E. O’Sullivan, “An update on PPAR activation by cannabinoids,” *British Journal of Pharmacology*, vol. 173, no. 12, pp. 1899–1910, 2016.
- [38] A. J. Hill, N. A. Jones, I. Smith et al., “Voltage-gated sodium (NaV) channel blockade by plant cannabinoids does not confer anticonvulsant effects per se,” *Neuroscience Letters*, vol. 566, pp. 269–274, 2014.
- [39] L. R. Ruhaak, J. Felth, P. C. Karlsson, J. J. Rafter, R. Verpoorte, and L. Bohlin, “Evaluation of the cyclooxygenase inhibiting effects of six major cannabinoids isolated from Cannabis sativa,” *Biological and Pharmaceutical Bulletin*, vol. 34, no. 5, pp. 774–778, 2011.
- [40] A. Alhamoruni, K. L. Wright, M. Larvin, and S. E. O’Sullivan, “Cannabinoids mediate opposing effects on inflammation-induced intestinal permeability,” *British Journal of Pharmacology*, vol. 165, no. 8, pp. 2598–2610, 2012.

- [41] D. G. Couch, H. Maudslay, B. Doleman, J. N. Lund, and S. E. O'Sullivan, "The use of cannabinoids in colitis: a systematic review and meta-analysis," *Inflammatory Bowel Diseases*, vol. 24, no. 4, pp. 680–697, 2018.
- [42] Y. Deguchi, A. Andoh, Y. Yagi et al., "The S1P receptor modulator FTY720 prevents the development of experimental colitis in mice," *Oncology Reports*, vol. 16, no. 4, pp. 699–703, 2006.
- [43] F. Borrelli, I. Fasolino, B. Romano et al., "Beneficial effect of the non-psychotropic plant cannabinoid cannabigerol on experimental inflammatory bowel disease," *Biochemical Pharmacology*, vol. 85, no. 9, pp. 1306–1316, 2013.
- [44] M. Rajesh, P. Mukhopadhyay, G. Haskó, L. Liaudet, K. Mackie, and P. Pacher, "Cannabinoid-1 receptor activation induces reactive oxygen species-dependent and -independent mitogen-activated protein kinase activation and cell death in human coronary artery endothelial cells," *British Journal of Pharmacology*, vol. 160, no. 3, pp. 688–700, 2010.
- [45] P. Mukhopadhyay, M. Rajesh, H. Pan et al., "Cannabinoid-2 receptor limits inflammation, oxidative/nitrosative stress, and cell death in nephropathy," *Free Radical Biology and Medicine*, vol. 48, no. 3, pp. 457–467, 2010.
- [46] A. L. Dawidowicz, M. Olszowy-Tomczyk, and R. Typek, "CBG, CBD, Δ^9 -THC, CBN, CBGA, CBDA and Δ^9 -THCA as antioxidant agents and their intervention abilities in antioxidant action," *Fitoterapia*, vol. 152, Article ID 104915, 2021.
- [47] S. Giacoppo, A. Gugliandolo, O. Trubiani et al., "Cannabinoid CB2 receptors are involved in the protection of RAW264.7 macrophages against the oxidative stress: an in vitro study," *European Journal of Histochemistry*, vol. 61, no. 1, p. 2749, 2017.
- [48] A. Gugliandolo, F. Pollastro, G. Grassi, P. Bramanti, and E. Mazzon, "In In vitro model of neuroinflammation: efficacy of cannabigerol, a non-psychoactive cannabinoid," *International Journal of Molecular Sciences*, vol. 19, no. 7, p. 1992, 2018.
- [49] S. H. Baek, S. Du Han, C. N. Yook, Y. C. Kim, and J. S. Kwak, "Synthesis and antitumor activity of cannabigerol," *Archives of Pharmacal Research*, vol. 19, no. 3, pp. 228–230, 1996.
- [50] S. H. Baek, Y. O. Kim, J. S. Kwag, K. E. Choi, W. Y. Jung, and D. S. Han, "Boron trifluoride etherate on silica-A modified Lewis acid reagent (VII). Antitumor activity of cannabigerol against human oral epitheloid carcinoma cells," *Archives of Pharmacal Research*, vol. 21, no. 3, pp. 353–356, 1998.
- [51] A. Ligresti, A. S. Moriello, K. Starowicz et al., "Antitumor activity of plant cannabinoids with emphasis on the effect of cannabidiol on human breast carcinoma," *Journal of Pharmacology and Experimental Therapeutics*, vol. 318, no. 3, pp. 1375–1387, 2006.
- [52] J. D. Wilkinson and E. M. Williamson, "Cannabinoids inhibit human keratinocyte proliferation through a non-CB1/CB2 mechanism and have a potential therapeutic value in the treatment of psoriasis," *Journal of Dermatological Science*, vol. 45, no. 2, pp. 87–92, 2007.
- [53] L. Beaugerie, "Inflammatory bowel disease therapies and cancer risk: where are we and where are we going?" *Gut*, vol. 61, no. 4, pp. 476–483, 2012.
- [54] F. Borrelli, E. Pagano, B. Romano et al., "Colon carcinogenesis is inhibited by the TRPM8 antagonist cannabigerol, a Cannabis-derived non-psychotropic cannabinoid," *Carcinogenesis*, vol. 35, no. 12, pp. 2787–2797, 2014.
- [55] T. T. Lah, M. Novak, M. A. Pena Almidon et al., "Cannabigerol is a potential therapeutic agent in a novel combined therapy for glioblastoma," *Cells*, vol. 10, no. 2, p. 340, 2021.
- [56] V. Di Giacomo, A. Chiavaroli, L. Recinella et al., "Antioxidant and neuroprotective effects induced by cannabidiol and cannabigerol in rat CTX-TNA2 astrocytes and isolated cortexes," *International Journal of Molecular Sciences*, vol. 21, no. 10, p. 3575, 2020.
- [57] N. L. Stone, T. J. England, and S. E. O'Sullivan, "Protective effects of cannabidivarin and cannabigerol on cells of the blood-brain barrier under ischemic conditions," *Cannabis and Cannabinoid Research*, vol. 6, no. 4, pp. 315–326, 2021.
- [58] S. Valdeolivas, C. Navarrete, I. Cantarero, M. L. Bellido, E. Muñoz, and O. Sagredo, "Neuroprotective properties of cannabigerol in Huntington's disease: studies in R6/2 mice and 3-nitropropionate-lesioned mice," *Neurotherapeutics*, vol. 12, no. 1, pp. 185–199, 2015.
- [59] A. G. Granja, F. Carrillo-Salinas, A. Pagani et al., "A cannabigerol quinone alleviates neuroinflammation in a chronic model of multiple sclerosis," *Journal of Neuroimmune Pharmacology*, vol. 7, no. 4, pp. 1002–1016, 2012.
- [60] E. S. Huseby, D. Kamimura, Y. Arima, C. S. Parello, K. Sasaki, and M. Murakami, "Role of T cell-glia cell interactions in creating and amplifying central nervous system inflammation and multiple sclerosis disease symptoms," *Frontiers in Cellular Neuroscience*, vol. 9, p. 295, 2015.
- [61] F. J. Carrillo-Salinas, C. Navarrete, M. Mecha et al., "A cannabigerol derivative suppresses immune responses and protects mice from experimental autoimmune encephalomyelitis," *PLoS One*, vol. 9, no. 4, Article ID e94733, 2014.
- [62] J. Díaz-Alonso, J. Paraíso-Luna, C. Navarrete et al., "VCE-003.2, a novel cannabigerol derivative, enhances neuronal progenitor cell survival and alleviates symptomatology in murine models of Huntington's disease," *Scientific Reports*, vol. 6, no. 1, Article ID 29789, 2016.
- [63] C. Rodríguez-Cueto, I. Santos-García, L. García-Toscano et al., "Neuroprotective effects of the cannabigerol quinone derivative VCE-003.2 in SOD1 (G93A) transgenic mice, an experimental model of amyotrophic lateral sclerosis," *Biochemical Pharmacology*, vol. 157, pp. 217–226, 2018.
- [64] C. García, M. Gómez-Cañas, S. Burgaz et al., "Benefits of VCE-003.2, a cannabigerol quinone derivative, against inflammation-driven neuronal deterioration in experimental Parkinson's disease: possible involvement of different binding sites at the PPAR γ receptor," *Journal of Neuroinflammation*, vol. 15, no. 1, p. 19, 2018.
- [65] S. Burgaz, C. García, M. Gómez-Cañas et al., "Neuroprotection with the cannabigerol quinone derivative VCE-003.2 and its analogs CBGA-Q and CBGA-Q-salt in Parkinson's disease using 6-hydroxydopamine-lesioned mice," *Molecular and Cellular Neuroscience*, vol. 110, Article ID 103583, 2021.
- [66] J. Aguarales, J. Paraíso-Luna, B. Palomares et al., "Oral administration of the cannabigerol derivative VCE-003.2 promotes subventricular zone neurogenesis and protects against mutant huntingtin-induced neurodegeneration," *Translational Neurodegeneration*, vol. 8, no. 1, 2019.
- [67] B. Lutz, "Endocannabinoid signals in the control of emotion," *Current Opinion in Pharmacology*, vol. 9, no. 1, pp. 46–52, 2009.
- [68] D. R. Janero and A. Makriyannis, "Cannabinoid receptor antagonists: pharmacological opportunities, clinical experience, and translational prognosis," *Expert Opinion on Emerging Drugs*, vol. 14, no. 1, pp. 43–65, 2009.
- [69] E. M. Rock, J. M. Goodwin, C. L. Limebeer et al., "Interaction between non-psychotropic cannabinoids in marijuana: effect of cannabigerol (CBG) on the anti-nausea or anti-emetic

- effects of cannabidiol (CBD) in rats and shrews,” *Psychopharmacology (Berl)*, vol. 215, no. 3, pp. 505–512, 2011.
- [70] J. De Vry, “5-HT_{1A} receptor agonists: recent developments and controversial issues,” *Psychopharmacology (Berl)*, vol. 121, no. 1, pp. 1–26, 1995.
- [71] L. D. O’Brien, K. L. Wills, B. Segsworth et al., “Effect of chronic exposure to rimonabant and phytocannabinoids on anxiety-like behavior and saccharin palatability,” *Pharmacology Biochemistry and Behavior*, vol. 103, no. 3, pp. 597–602, 2013.
- [72] A. Zagzoog, K. A. Mohamed, H. J. J. Kim et al., “In vitro and in vivo pharmacological activity of minor cannabinoids isolated from *Cannabis sativa*,” *Scientific Reports*, vol. 10, no. 1, Article ID 20405, 2020.
- [73] C. L. Limebeer and L. A. Parker, “The 5-HT_{1A} agonist 8-OH-DPAT dose-dependently interferes with the establishment and the expression of lithium-induced conditioned rejection reactions in rats,” *Psychopharmacology (Berl)*, vol. 166, no. 2, pp. 120–126, 2003.
- [74] J. A. Galli, R. Andari Sawaya, and F. K. Friedenberg, “Cannabinoid hyperemesis syndrome,” *Current Drug Abuse Reviews*, vol. 4, no. 4, pp. 241–249, 2011.
- [75] R. S. Hepler and I. R. Frank, “Marihuana smoking and intraocular pressure,” *JAMA*, vol. 217, no. 10, p. 1392, 1971.
- [76] W. D. Purnell and J. M. Gregg, “Delta (9)-tetrahydrocannabinol, euphoria and intraocular pressure in man,” *Annals of Ophthalmology*, vol. 7, no. 7, pp. 921–923, 1975.
- [77] B. K. Colasanti, C. R. Craig, and R. D. Allara, “Intraocular pressure, ocular toxicity and neurotoxicity after administration of cannabiniol or cannabigerol,” *Experimental Eye Research*, vol. 39, no. 3, pp. 251–259, 1984.
- [78] B. K. Colasanti, “A comparison of the ocular and central effects of delta 9-tetrahydrocannabinol and cannabigerol,” *Journal of Ocular Pharmacology and Therapeutics*, vol. 6, no. 4, pp. 259–269, 1990.
- [79] A. M. Szczesniak, Y. Maor, H. Robertson, O. Hung, and M. E. Kelly, “Nonpsychotropic cannabinoids, abnormal cannabidiol and cannabigerol-dimethyl heptyl, act at novel cannabinoid receptors to reduce intraocular pressure,” *Journal of Ocular Pharmacology and Therapeutics*, vol. 27, no. 5, pp. 427–435, 2011.
- [80] M. Pucci, C. Rapino, A. Di Francesco, E. Dainese, C. D’Addario, and M. Maccarrone, “Epigenetic control of skin differentiation genes by phytocannabinoids,” *British Journal of Pharmacology*, vol. 170, no. 3, pp. 581–591, 2013.
- [81] L. K. Dunn, J. L. O’Neill, and S. R. Feldman, “Acne in adolescents: quality of life, self-esteem, mood, and psychological disorders,” *Dermatology Online Journal*, vol. 17, no. 1, p. 1, 2011.
- [82] A. Oláh, A. Markovics, J. Szabó-Papp et al., “Differential effectiveness of selected non-psychotropic phytocannabinoids on human sebocyte functions implicates their introduction in dry/seborrhoeic skin and acne treatment,” *Experimental Dermatology*, vol. 25, no. 9, pp. 701–707, 2016.
- [83] E. Pagano, V. Montanaro, A. Di Girolamo et al., “Effect of non-psychotropic plant-derived cannabinoids on bladder contractility: focus on cannabigerol,” *Natural Product Communications*, vol. 10, no. 6, p. 1012, 2015.
- [84] M. R. Ruggieri, “Cannabinoids: potential targets for bladder dysfunction,” *Handbook of Experimental Pharmacology*, vol. 202, pp. 425–451, 2011.
- [85] E. A. Formukong, A. T. Evans, and F. J. Evans, “The inhibitory effects of cannabinoids, the active constituents of *Cannabis sativa* L. on human and rabbit platelet aggregation,” *Journal of Pharmacy and Pharmacology*, vol. 41, no. 10, pp. 705–709, 2011.
- [86] G. Appendino, S. Gibbons, A. Giana et al., “Antibacterial cannabinoids from *Cannabis sativa*: a structure-activity study,” *Journal of Natural Products*, vol. 71, no. 8, pp. 1427–1430, 2008.
- [87] M. Aqawi, R. V. Sionov, R. Gallily, M. Friedman, and D. Steinberg, “Anti-bacterial properties of cannabigerol toward *Streptococcus mutans*,” *Frontiers in Microbiology*, vol. 12, Article ID 656471, 2021.
- [88] M. Aqawi, R. Gallily, R. V. Sionov, B. Zaks, M. Friedman, and D. Steinberg, “Cannabigerol prevents quorum sensing and biofilm formation of *Vibrio harveyi*,” *Frontiers in Microbiology*, vol. 11, p. 858, 2020.
- [89] D. I. Brierley, J. Samuels, M. Duncan, B. J. Whalley, and C. M. Williams, “Cannabigerol is a novel, well-tolerated appetite stimulant in pre-satiated rats,” *Psychopharmacology (Berl)*, vol. 233, no. 19–20, pp. 3603–3613, 2016.
- [90] D. I. Brierley, J. Samuels, M. Duncan, B. J. Whalley, and C. M. Williams, “A cannabigerol-rich *Cannabis sativa* extract, devoid of Δ^9 -tetrahydrocannabinol, elicits hyperphagia in rats,” *Behavioural Pharmacology*, vol. 28, no. 4, pp. 280–284, 2017.
- [91] D. I. Brierley, J. R. Harman, N. Giallourou et al., “Chemotherapy-induced cachexia dysregulates hypothalamic and systemic lipamines and is attenuated by cannabigerol,” *Journal of Cachexia, Sarcopenia and Muscle*, vol. 10, no. 4, pp. 844–859, 2019.
- [92] N. M. Kogan, Y. Lavi, L. M. Topping et al., “Novel CBG derivatives can reduce inflammation, pain and obesity,” *Molecules*, vol. 26, no. 18, p. 5601, 2021.