

AN INTRONIC VARIANT IN *PTEN* GENE IS PROBABLY INVOLVED IN SPLICING ALTERATION EVENTS.

Carmela Rinaldi, José Freni

Department of Biomedical and Dental Sciences and Morphofunctional Imaging, Division of Medical Biotechnologies and Preventive Medicine, University of Messina, Messina, Italy

ARTICLE INFO

Article history:

Received 19 January 2019

Revised 19 April 2019

Accepted 11 May 2019

Keywords:

PTEN, tumor suppressor, breast cancer.

ABSTRACT

PTEN is a gate-keeper tumor-suppressor gene involved in various cell cycle pathways. Germline mutations of one allele of *PTEN* were found associated with Cowden syndrome, which is an inherited disease characterized by multiple hamartomas. *PTEN* is frequently inactivated also by somatic mutations in various array of human tumors. In this study, a cohort of 50 Sicilian breast cancer patients, negative to screening for pathogenic mutations in *BRCA1/2* genes, was analyzed. Variant c.492+46 G>A (rs775537003) in intron 5 of *PTEN* gene was identified. The possible role of this variant in splicing mechanisms was evaluated by in silico analysis. Alteration of an ESE and ESS site was found. This is a preliminary study with the aim to investigate about the involvement of *PTEN* gene in the breast cancer and the possible role of splicing alteration events into the risk of the disease.

© EuroMediterranean Biomedical Journal 2019

1. Introduction

PTEN (phosphatase and tensin homolog deleted on chromosome 10) is a tumor-suppressor gene which encodes a major lipid 3-phosphatase. This protein signals down the PI3 kinase/AKT proapoptotic pathway and various cell-survival pathways, such as the mitogen-activated kinase (MAPK) pathway. The role of *PTEN* mainly consists in the ability to dephosphorylate and degrade PIP3 phospholipid of membrane, and consequently inhibit serine/threonine kinase oncogenic AKT [1]. Furthermore, *PTEN* coordinates G1 arrest through up-regulation of p27, an inhibitor of cyclin-dependent kinases, and acts as a negative regulator of the cell cycle and concomitant down-regulation of cyclin D1 [2]. Originally germline mutations of one allele of *PTEN* were found associated with Cowden syndrome (CS), which is an inherited disease characterized also by hamartomatous gastrointestinal polyposis that can evolve into early-onset esophagogastric cancer. [3-5].

PTEN is frequently inactivated by somatic mutations in a wide array of human tumors. Selective inactivation of *PTEN* in the developing brain causes defects in the neuronal migration, synapses formation and stabilization, and progression of glioblastoma multiforme (GBM), one the most malignant astrocytic tumor [6]. The astrocytes play an important role in several different biologic activities and the *PTEN* loss can increase the proliferation of astrocytes in vivo [7].

Previous studies have shown that glial fibrillary acidic protein (GFAP) co-localizes with a wide range of proteins expressed in astrocytes, as sarcoglycans and dystrophin complex. The loss of some elements of these complex could be cause of the loss of structural plasticity of the synapses resulting in a tumor progression.

Other neoplastic diseases in which *PTEN* takes part is prostate cancer; however, in this case, mechanism involves the glyoxalases (GLOs) [8], a class of enzymes involved in cellular detoxification from Reactive Oxygen Species (ROS); moreover, polymorphisms in GLO genes were associated to several pathological conditions affecting Central Nervous System and retina [9-11]. Their role in breast cancer is, instead, controversial [12-13]. Importantly, the reduced expression of *PTEN* is not only due to loss of heterozygosity (LOH) but also to epigenetic mechanisms, as methylation of promoter [14-15]. This phenomenon was also observed in endothelial cells derived from Cerebral Cavernous Malformations (CCMs) sporadic lesions [16]. CCMs are benign neoplasms derived from angiogenic impairment due to mutations at three loci [17-19]. *PTEN*/VEGF signaling modulates the three CCM proteins activity [20].

In present study on a cohort of 50 Sicilian breast cancer patients, negative to screening for pathogenic mutations in *BRCA1/2* genes, the entire coding sequence of *PTEN* gene was analyzed in order to evaluate the role of this gene in the disease.

2. Methods

Patients

The peripheral blood samples were collected from 50 women (over 50 years of age) living in Sicily. Before recruitment, each patient was asked to sign an informed consent form. No adjuvant chemotherapy was administered. We have no information on treatment with radiation and/or endocrine therapy.

The study followed the tenets of the Declaration of Helsinki and was approved by the Scientific Ethics Committee of the Azienda Ospedaliera Universitaria-Policlinico "G. Martino" Messina.

Molecular analysis

Primers and polymerase chain reaction (PCR). The heparinized peripheral blood samples were collected from breast cancer patients. Genomic DNA was extracted from lymphocytes by kit QI Amp DNA Blood Mini Kit (QIAGEN®).

The nucleotide sequence of primers employed for PCR reaction is shown in Table 1.

Gene	Exon	Sequence primer	Amplicon size (bp)	Ta
PTEN	1	PF: 5'-AAGTCCAGAGCCATTTCCATC-3'	261	59,5°C
		PR: 5'-AACTACGGACATTTTCGCATC-3'		
PTEN	2	PF: 5'-AGTTTGATTGCTGCATATTTTC-3'	300	56,5°C
		PR: 5'-GAAGTCCATTAGGTACGGTAA-3'		
PTEN	3	PF: 5'-TGTAATTTCAAATGTTAGCTC-3'	258	53,5°C
		PR: 5'-TCTTGACTTCTTGACTTAAT-3'		
PTEN	4	PF: 5'-GTGATAACAGTATCTACTTAAT-3'	284	54,5°C
		PR: 5'-TGACAGTAAGATACAGCTAT-3'		
PTEN	5	PF: 5'-CTGTTAAGTTTGTATGCAACA-3'	451	54,5°C
		PR: 5'-ACTTGTCAATTACACCTCAAT-3'		
PTEN	6	PF: 5'-CCAGTTACCATAGCAATTTAGT-3'	266	58,5°C
		PR: 5'-CTGTTCCAATACATGGAAGGA-3'		
PTEN	7	PF: 5'-ATTGACAGATACAGAATCCATA-3'	410	55,5°C
		PR: 5'-AAA ACA CCT GCA GAT CTA ATA-3'		
PTEN	8	PF: 5'-AATGTTTAAACATAGGTGACAG-3'	426	53,5°C
		PR: 5'-TTGCAAGCAAGTTCCTCAT-3'		
PTEN	9	PF: 5'-AAGATCATGTTTGTACAGTG-3'	428	55,5°C
		PR: 5'-ATCTGACACAATGTCCTATTG-3'		

PF= forward; PR=reverse; Ta=annealing temperature.

Table 1. Primer sequences used for PCR and sequencing reaction

PCR reaction was carried out adding 0.8 µg of genomic DNA to 50 µl reaction mixture containing a 0.3 µM concentration of each primer and 1.5 U MyTaq polymerase (Bioline). It was carried out in the thermal cycler (Gene Amp PCR System 2700; PE Applied Biosystems, Foster City, CA) under following conditions: denaturation at 95°C for 15 sec, annealing for 15 sec and extension at 72°C for 15 sec for 35 cycles, after an initial 1 min denaturation at 95 °C. The final extension step at 72°C was extended to 10 minutes. Annealing temperature was optimized for each primer set as described in Table 1.

The PCR products were resolved on 1% agarose gel and visualized by ethidium bromide staining.

Sequencing was carried out on the primary amplicons purified by kit "ISOLATE II PCR and Gel Kit (BIOLINE)", using the DNA direct sequencing with the ABI PRISM® Big Dye® Terminator v3.1 Cycle Sequencing Kit on the capillary electrophoresis analyzer 3500 Dx System (Applied Biosystems).

The specificity for the detected variants was tested by NCBI GeneBank, <http://www.ncbi.nlm.nih.gov/genbank/> and NCBI Blast data base, <http://blast.ncbi.nlm.nih.gov/Blast.cgi>.

In silico analysis

Intron variant c.492+46 G>A was found and its possible role in splicing mechanisms was evaluated. The possible creation of exonic splicing silencer (ESS) and the abolition of exonic splicing enhancer (ESE) sites, as well as the abolition and creation of canonical sites and cryptic splice sites respectively, were analyzed through an in silico analysis using Human Splicing Finder software (<http://www.umd.be/HSF3/HSF.html>) [21].

3. Results

The mutation analysis was performed in the patients negative for pathogenic mutations in BRCA1/2 genes. The analysis of the entire coding sequence of PTEN gene, performed on 9 exons and partial intron sequences upstream and downstream of each exon (+/- nucleotides) identified the variant c.492+46 G>A (rs775537003) (http://www.ensembl.org/Homo_sapiens/Gene/Sequence?db=core;g=ENS G00000171862;r=10:87863113-87971930) in intron 5, a few dozen nucleotides upstream of exon 6.

Although this variant does not alter the structure of the protein, it could be involved in splicing mechanism. To evaluate the possible role of the variant in the splicing mechanism, in silico analysis was performed by Human Splicing Finder. The software revealed the alteration of an exonic splicing enhancer (ESE) site and an exonic splicing silencer (ESS) site. (Figure 1).

Sequence Position	cDNA Position	Enhancer motif reference sequence	Enhancer motif mutant sequence	Variation
402	+402		ctgaaa	New Site
403	+403	tgagaa		Site broken
404	+404		gaaaat	New Site

Sequence Position	cDNA Position	Silencer motif reference sequence	Silencer motif mutant sequence	Variation
406	+406	gaattt		Site broken

Figure 1. Possible alteration of the splicing process due to intronic c.492+46 G>A variant.

In addition, a reduction in SF2/ASF pattern score may be correlated with an exon-skipping phenotype associated with various disease-causing mutations, such as breast cancer (doi: 10.1093/hmg/ddl171) (Figure 2).

Sequence Position	cDNA Position	Linked SR protein	Reference Motif (value 0-100)	Linked SR protein	Mutant Motif (value 0-100)	Variation
399	+399	SC35	galcigag (77.75)			Site broken -100
402	+402	SF2/ASF (IgM-BRCA1)	ctgagaa (76.23)			Site broken -100
402	+402	SF2/ASF	ctgagaa (73.50)			Site broken -100

Figure 2 - Reduction in SF2/ASF pattern score due to c.492+46 G>A variant.

4. Discussion

More than 50 genes are involved in the progression of breast cancer. Over 1500 variants in two principal genes (BRCA1 and BRCA2) were listed in the Breast Cancer Information Core (BIC) [22-23].

In this study, the choice about *PTEN* is first linked to the important pathway largely controlled by this gene, to its role as gate-keeper tumor-suppressor gene, but above all because it is an haploinsufficient tumor-suppressor gene. Therefore, a single mutation can determine the production of a dysfunctional phosphatase.

Furthermore, the variant identified in *PTEN* gene is a polymorphism. To establish potential pathogenic or protective role of polymorphisms in diseases' development is very complex; while, for rare disease, few cases may be sufficient [24-27], for breast cancer, having a wide incidence, a very large cohort of patients is required. Moreover, this study was limited to *PTEN* coding regions and intron/exon boundaries and did not consider possibility of presence of variants in the promoter region that was shown may be interfere with normal gene expression [28-30].

Certainly, further functional studies will be necessary to confirm the lack of activity of the aberrant protein form. Other unidentified genes should be screened in order to complete the framework in the field of the breast cancer.

References

- Sun H, Lesche R, Li DM, Liliental J, Zhang H, Gao J, Gavrilova N, Mueller B, Liu X, Wu H: PTEN modulates cell cycle progression and cell survival by regulating phosphatidylinositol 3,4,5-triphosphate and Akt/protein kinase B signaling pathway. *Proc Natl Acad Sci U S A* 1999; 96:6199-6204.
- Rinaldi C, Malara NM, D'Angelo R, Sidoti A, Leotta A, Lio S, Caparello B, Ruggeri A, Mollace V, Amato A: Age dependent switching role of cyclin D1 in breast cancer. *Analytical Cellular Pathology* 2012;35:179-185.
- Singh B, Ittmann MM, Krolewski JJ: Sporadic breast cancers exhibit loss of heterozygosity on chromosome segment 10q23 close to the Cowden disease locus. *Genes Chromosomes Cancer* 1998; 21:166-171.
- Saraniti C, Gallina S, Santangelo M, Montana F, Portelli F, Orlando E: Sinonasal respiratory epithelial adenomatoid hamartoma: a report on three cases. *EuroMediterranean Biomedical Journal* 2018; 13:060-064

- Milazzotto R, Spatola C, Tocco A, Raffaele L, Salamone V, Aglieco A, Trombatore C, Di Cataldo A, Privitera G: Metastatic esophagogastric junction cancer: case report of a complete long-term response after combined chemoradiotherapy. *EuroMediterranean Biomedical Journal* 2018; 13:082-084
- Lachyankar MB, Sultana N, Schonhoff CM, Mitra P, Poluha W, Lambert S, Quesenberry PJ, Litofsky NS, Recht LD, Nabi R, Miller SJ, Ohta S, Neel BG, Ross AH: A role for nuclear PTEN in neuronal differentiation. *J Neurosci* 2000; 20:1404-1413.
- Fraser MM, Zhu X, Kwon CH, Uhlmann EJ, Gutmann DH, Baker SJ: Pten loss causes hypertrophy and increased proliferation of astrocytes in vivo. *Cancer Res* 2004; 64:7773-7779.
- Valenti D, Vacca RA, de Bari L: 3-Bromopyruvate induces rapid human prostate cancer cell death by affecting cell energy metabolism, GSH pool and the glyoxalase system. *J Bioenerg Biomembr* 2015;47:493-506.
- Rinaldi C, Bramanti P, Famà A, Scimone C, Donato L, Antognelli C, Alafaci C, Tomasello F, D'Angelo R, Sidoti A: Glyoxalase I A111E, Paraoxonase 1 Q192R and L55M polymorphisms in Italian patients with sporadic cerebral cavernous malformations; a pilot study. *J Biol Regul Homeost Agents* 2015;29:493-500.
- Tao H, Si L, Zhou X, Liu Z, Ma Z, Zhou H, Zhong W, Cui L, Zhang S, Li Y, Ma G, Zhao J, Huang W, Yao L, Xu Z, Zhao B, Li K: Role of glyoxalase I gene polymorphisms in late-onset epilepsy and drug-resistant epilepsy. *J Neurol Sci* 2016; 363:200-206.
- Donato L, Scimone C, Nicocia G, Denaro L, Robledo R, Sidoti A, D'Angelo R: GLO1 gene polymorphisms and their association with retinitis pigmentosa: a case-control study in a Sicilian population. *Mol Biol Rep* 2018;45:1349-1355.
- Germanová A, Germanová A, Tesarová P, Jáchymová M, Zvára K, Zima T, Kalousová M: Glyoxalase I Glu111Ala polymorphism in patients with breast cancer. *Cancer Invest* 2009;27:655-660.
- Naidu R, Har YC, Taib NA: Glyoxalase I Ala111Glu gene polymorphism: No association with breast cancer risk but correlated with absence of progesterone receptor. *Pathol Int* 2010;60:614-620.
- Baeza N, Weller M, Yonekawa Kleihues, P, Ohgaki, H: PTEN methylation and expression in glioblastomas. *Acta Neuropathol (Berlin)* 2003;106:479-485.
- Garcia JM, Silva J, Pena C, Garcia V, Rodriguez R, Cruz MA, Cantos B, Provencio M, Espana P, Bonilla F: Promoter methylation of the PTEN gene is a common molecular change in breast cancer. *Genes Chromosomes Cancer* 2004; 41:117-124.
- Zhu Y, Wloch A, Wu Q, Peters C, Pagenstecher A, Bertalanffy H, Sure U: Involvement of PTEN promoter methylation in cerebral cavernous malformations. *Stroke* 2009;40:820-826.
- D'Angelo R, Scimone C, Calabrò M, Schettino C, Fratta M, Sidoti A: Identification of a novel CCM2 gene mutation in an Italian family with multiple cerebral cavernous malformations and epilepsy: a causative mutation? *Gene* 2013; 519:202-207.
- Scimone C, Bramanti P, Ruggeri A, Katsarou Z, Donato L, Sidoti A, D'Angelo R: Detection of Novel Mutation in Ccm3 Causes Familial Cerebral Cavernous Malformations. *J Mol Neurosci* 2015;57:400-403.

19. Scimone C, Bramanti P, Alafaci C, Granata F, Piva F, Rinaldi C, Donato L, Greco F, Sidoti A, D'Angelo R: Update on Novel CCM Gene Mutations in Patients with Cerebral Cavernous Malformations. *J Mol Neurosci* 2017; 61:189-198.
20. Kar S, Samii A, Bertalanffy H: PTEN/PI3K/Akt/VEGF signaling and the cross talk to KRIT1, CCM2, and PDCD10 proteins in cerebral cavernous malformations. *Neurosurg Rev* 2015; 38:229-236.
21. Desmet FO, Hamroun D, Lalande M, Collod-B  roud G, Claustres M, B  roud C: Human Splicing Finder:an online bioinformatics tool to predict splicing signals. *Nucleic Acids Res* 2009; 37,e67.
22. Szabo C, Masiello A, Ryan JF, Brody LC: The breast cancer information core: database design, structure, and scope. *Hum Mutat* 2000;16:123-131.
23. Couch FJ, Nathanson KL, Offit K: Two decades after BRCA: setting paradigms in personalized cancer care and prevention. *Science* 2014;343,1466-1470.
24. D'Angelo R, Scimone C, Rinaldi C, Trimarchi G, Italiano D, Bramanti P, Amato A, Sidoti A: CCM2 gene polymorphisms in Italian sporadic patients with cerebral cavernous malformation: a case-control study. *Int J Mol Med* 2012;29:1113-1120.
25. D'Angelo R, Donato L, Venza I, Scimone C, Aragona P, Sidoti A: Possible protective role of the ABCA4 gene c.1268A>G missense variant in Stargardt disease and syndromic retinitis pigmentosa in a Sicilian family: Preliminary data. *Int J Mol Med* 2017;39:1011-1020.
26. Donato L, Bramanti P, Scimone C, Rinaldi C, D'Angelo R, Sidoti A: miRNA expression profile of retinal pigment epithelial cells under oxidative stress conditions. *FEBS Open Bio* 2018; 8:219-233.
27. Robledo R, Orru S, Sidoti A, Muresu R, Esposito D, Grimaldi MC, Carcassi C, Rinaldi A, Bernini L, Contu L, Romani M, Roe B, Siniscalco M: A 9.1-kb gap in the genome reference map is shown to be a stable deletion/insertion polymorphism of ancestral origin. *Genomics* 2002; 80:585-592.
28. Scimone C, Bramanti P, Ruggeri A, Donato L, Alafaci C, Crisafulli C, Mucciardi M, Rinaldi C, Sidoti A, D'Angelo R: CCM3/SERPINI1 bidirectional promoter variants in patients with cerebral cavernous malformations: a molecular and functional study. *BMC Med Genet* 2016; 17:74.
29. Donato L, Scimone C, Rinaldi C, Aragona P, Briuglia S, D'Ascola A, D'Angelo R, Sidoti A: Stargardt Phenotype Associated With Two ELOVL4 Promoter Variants and ELOVL4 Downregulation: New Possible Perspective to Etiopathogenesis? *Invest Ophthalmol Vis Sci* 2018;59:843-857.
30. Donato L, Scimone C, Rinaldi C, D'Angelo R, Sidoti A: Non-coding RNAome of RPE cells under oxidative stress suggests unknown regulative aspects of Retinitis pigmentosa etiopathogenesis. *Sci Rep.* 2018;8:16638.