

A unique fatal case of Waterhouse–Friderichsen syndrome caused by *Proteus mirabilis* in an immunocompetent subject

Case report and literature analysis

Elvira Ventura Spagnolo, MD^{a,*}, Cristina Mondello, MD^b, Salvatore Rocuzzo, MD^b, Chiara Stassi^a, Luigi Cardia, MD^c, Angela Grieco, MD^d, Cataldo Raffino, MD^d

Abstract

Introduction: The Waterhouse–Friderichsen syndrome (WFS), also known as purpura fulminans, is a potentially lethal condition described as acute hemorrhagic necrosis of the adrenal glands. It is often caused by infection. Classically, *Neisseriae meningitidis* represents the main microorganism related to WFS, although, infrequently, also other infectious agents are reported as a possible etiologic agent. The authors report the first case of death due to *Proteus mirabilis* infection, with postmortem evidence of WFS.

Patient concerns: After a facial trauma that provoked a wound on the nose, the subject, a healthy 40-years old man, was conducted to the local hospital (in Sicily, Italy) after the primary care he was discharged. Subsequently, after 2 days of general malaise, he returned to the hospital due to the worsening of the clinical condition. During the hospitalization, hypotension, and neurological impairment appeared; the laboratory analysis showed leukocytosis and the alteration of renal, hepatic and coagulative parameters. Microbiological blood analysis resulted positive for a *P mirabilis* infection.

Diagnosis: Multiorgan failure (MOF) with disseminated intravascular coagulation (DIC) due to sepsis was diagnosed.

Interventions: The practitioners administered intensive support, antibiotic therapy, antithrombin III, vitamin K, and plasma.

Outcomes: After 3 days the subject died. The autopsy and the microscopic investigation were performed revealing, also, the adrenal diffuse micronodular hyperplasia associated with a cortico-medullary hemorrhagic apoplexy.

Conclusion: To our knowledge, this is the first case of MOF with WFS due to *P mirabilis* infection. This case report suggests that *P mirabilis* should be added to the list of unusual bacteria causing WFS. Furthermore, it supports the theory that any bacterium which causes DIC may cause adrenal hemorrhage and should suggest to clinicians the importance to consider a potential adrenal involvement in every patient with sepsis and DIC.

Abbreviations: ACTH = adrenocorticotrophic hormone, CT = computed tomography, DIC = disseminated intravascular dissemination, MOF = multiorgan failure, WFS = Waterhouse–Friderichsen syndrome.

Keywords: bilateral massive adrenal hemorrhage, forensic sciences, *Proteus mirabilis* infection, sepsis, Waterhouse-Friderichsen syndrome

Editor: N/A.

In the reported case there are no personal-specific medical information that allow to identify the subject because the described clinical picture, demonstrating septic shock, is common in many patients and the data concerning the patient have been sufficiently anonymized (eg, no name or photographs of the patient are included in the paper). Moreover, even if the patient is dead, we sought permission from a relative as a matter of courtesy and medical ethics. Unfortunately, no relatives were contactable. Finally, the authors got the permission to publish the case by the Public Prosecutor who performed the forensic investigations. Informed consent to publish the case report was obtained from the public prosecutor who performed the forensic investigations.

We asked for ethical approval but it was not necessary because the present work is a descriptive case report with no ethical implications.

The autopsy was performed for legal reasons because it was necessary to exclude the possibility that death was related to a crime.

The authors have no conflicts of interest to disclose.

^a Legal Medicine Section – Department for Health Promotion and Mother-Child Care, University of Palermo, Via del Vespro, ^b Department of Biomedical and Dental Sciences and Morphofunctional Imaging, University of Messina, ^c Department of Human Pathology of Adult and Childhood “Gaetano Barresi,” University of Messina, Messina, ^d Legal Medicine Centre of INAIL, Enna, Italy.

* Correspondence: Elvira Ventura Spagnolo, Department for Health Promotion and Mother-Child Care, University of Palermo, Via del Vespro, 129, 90127, Italy (e-mail: elvira.ventura@unipa.it).

Copyright © 2019 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Medicine (2019) 98:34(e16664)

Received: 24 January 2019 / Received in final form: 26 May 2019 / Accepted: 8 July 2019

<http://dx.doi.org/10.1097/MD.00000000000016664>

1. Introduction

The Waterhouse–Friderichsen syndrome (WFS), also known as purpura fulminans, is a very uncommon event but with high morbidity and mortality, that can occur in association with a common infection. It is described as acute hemorrhagic necrosis of the adrenal glands, often caused by meningococcal infection and is more frequently observed in the pediatric than the adult population.^[1]

Adrenal hemorrhage is an infrequent condition that can be associated both to noninfection (including postoperative state, thromboembolic diseases, anticoagulant treatment, burns, trauma, tumor metastasis, and cardiovascular catastrophes) and a variety of infectious diseases. Classically, *Neisseria meningitidis* represents the main microorganism related to WFS but is reported that also other infectious agents can be the possible etiologic agent of this disease, including bacteria (gram-negative and gram-positive) and viruses. Although *N meningitidis* accounts for more than 80% of WFS cases, other microorganisms implicated include other bacteria as *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, group A *Streptococcus*, *Neisseria gonorrhoeae*.^[2]

The authors report a case of bilateral massive adrenal hemorrhage due to bacteremia of *Proteus mirabilis* that, according to the literature, is the first case of WFS occurring in the setting of this bacterial infection. Also, the authors report an overview of the syndrome and the main uncommon organisms that are related to this disease.

2. Case report

In winter 2017, a 40 years-old Caucasian man was conducted to the emergency department of the local hospital in Sicily (Italy) after a facial trauma that caused a wound on the nose. On physical examination, no abnormal findings were found, and cranium computed tomography (CT) was negative. The wound was sutured and the patient was discharged. After 2 days, the subject returned to the hospital for a state of unspecific general malaise, and then was transferred to the emergency department where the clinicians observed a neurological impairment (Glasgow Coma Scale: 10) state. Laboratory investigations showed severe hypoglycemia (18 mg/dL), a high anion gap metabolic acidosis; and a glucose-bicarbonate solution was infused. After a few hours, the clinical picture worsened: the man was appearing confused, with the blood pressure of 80/40 mm Hg, heart rate of 96 b/min, oxygen saturation of 96% (with oxygen administration via Ventimask). The other laboratory analyses have shown renal and hepatic impairment, severe thrombocytopenia (5×10^3 /mL platelets), partial thromboplastin time 42.2 seconds, prothrombin time 17.2 seconds, prothrombin activity 56%, international normalized ratio 1.46, leukocytosis (29×10^3 /mL white blood cells). These data were indicative for severe multiorgan failure (MOF) with disseminated intravascular coagulation (DIC). Abdomen-chest CT revealed a diffuse swelling of both adrenal glands and inflammatory changes in the peri-adrenal and renal fat. The patient was transferred to the intensive care unit and then was intubated; the administration of antibiotic therapy (meropenem, 1 g per day iv), antithrombin III, vitamin K, dobutamine, noradrenalin, and plasma was started. During the following days, the patient remained in critical conditions and the blood microbiological analysis showed positivity for *P mirabilis* (meropenem sensible $<0.25 \mu\text{g/mL}$). In the night of the sixth day of hospitalization, the patient was

suddenly nonstable hemodynamically and, despite the administration of noradrenaline and adrenaline over the range, he developed a status of severe shock and died. The Judicial Authority ordered the autopsy. The autopsy showed brain congestion and pulmonary hemorrhagic oedema. Splenomegaly, hyperplasia of mesenteric lymph nodes, erosive and hemorrhagic lesions in the mucosa of stomach and intestine were observed. The adrenal lodges showed a fibrino-hemorrhagic infiltration (Fig. 1A and B).

The histological analysis showed endobronchial-endoalveolar haemorrhages and arteriolar lung embolus. Spleen had hemorrhagic congestion of the red pulp and hyperplasia of the white pulp. Intestine showed erosive lymphomonocytic inflammation, cytoateolonecrosis lesions of the submucosa and lymphomonocytic inflammation of the myenteric plexus; in the large intestine were also observed submucosal foci of micro-thromboembolism. Both adrenal glands showed diffuse micronodular hyperplasia associated with a cortico-medullary hemorrhagic apoplexy (Fig. 1C).

3. Discussion and conclusions

Infections and septic shock are among the most common causes of death that, not infrequently, are related to sudden unexpected death and medical malpractice.^[3–5] The present case, in fact, initially come to authors attention to evaluate if there were any professional liability profiles, that, on the basis of the cause of death and management of the patient, that were subsequently excluded. The clinical data and, in particular, the autopsy and histological findings, suggested that the subject died for MOF associated with DIC and bilateral adrenal apoplexy (WFS) due to sepsis sustained by *P mirabilis*. The clinical danger entailed in bilateral adrenal hemorrhage is acute adrenal insufficiency in the setting of severe illness, which can lead to rapid cardiovascular collapse and death.^[6] The acute hemorrhagic necrosis of the adrenal glands is considered a subtype of severe infection or endotoxemia and a lot of microorganisms that can induce DIC can lead to intravascular changes of the adrenal vessels (forming a unique blood supply, where an arterial subcapsular plexus drains on relatively few venules) and to apoplexy adrenal glands. Thus, endotoxin-mediated septic shock is the most frequent clinical condition seen in patients with infectious adrenal hemorrhage. These endotoxins (lipopolysaccharide in the case of Gram-negative bacteria and peptidoglycan for Gram-positive bacteria), bind the toll-like receptors of endothelial and inflammatory cells, inducing a number of signaling pathways that lead to transcription of coagulation, fibrinolysis, and pro-inflammatory cytokine genes activation (as tissue factor, interleukin [IL]-1, IL-6, and tumor necrosis factor-alpha).^[7]

It has been proposed that uni-visceral (or single organ type) Shwartzman mechanism may play an important role in the pathogenesis of adrenal haemorrhage and necrosis in humans. Since stress induces adrenal over-function through adrenocorticotrophic hormone (ACTH) secretion, it can be claimed that stress is capable to injure the adrenal tissue directly, or that the shock state, caused by successive stresses, may provoke peripheral circulatory disturbance which would influence the adrenal gland.^[8] The ACTH stimulates cortisol synthesis and secretion by regulating multiple steps in the steroidogenetic pathway. Activation of the melanocortin receptor 2 by ACTH in the adrenals also induces the adrenal production of factors affecting adrenal growth and its blood flow. Furthermore, ACTH




Figure 1. (A and B) Gross examination of adrenal nodules showing the presence of bilateral (respectively A: right; B: left) fibrino-hemorrhagic deposition; (C) Microscopic examination of adrenal gland showing diffuse micronodular hyperplasia associated with cortico-medullary hemorrhagic apoplexy.

stimulates the intra-adrenal production of vascular endothelial growth factor and the vaso-relaxant epoxy-eicosatrienoic acids.^[9,10]

However, it is also described that diffuse haemorrhage and necrosis of the adrenal do not occur easily either after ACTH administration or under stress conditions in humans. Additionally, stronger stimulation may be necessary to provoke a real lesion of the adrenal, and this could be given by an infection or by endotoxemia.^[8]

Concerning the triggering factor represented by endotoxemia, even if the *N meningitidis* is the most common pathogens associated with WFS, other microorganisms were identified as the possible cause of this adrenal disease. These microorganisms are *H influenzae*, *S pneumoniae*, *S. aureus*, group A *Streptococcus*, and *N gonorrhoeae*. The literature shows, also, unique or rare cases in which the WFS was related to infection caused by *Pseudomonas aeruginosa*, *Klebsiella oxytoca*, *Ewingella americana*, *Capnocytophaga canimorsus*, *Escherichia coli*, *Acinetobacter calcoaceticus*, DF-2 (Dysonian fermenter 2) bacterium, *Pasteurella multocida*, *Moraxella duplex*, *Rickettsia rickettsia*, *Bacillus anthracis*, *Ehrlichia* sp., *Treponema pallidum*, *Legionella pneumophila*, *Plesiomonas shigelloides* and also by viruses as Varicella zoster.^[2,11–20] The main information about these cases is summarized in Table 1.

Patients with the WFS usually show a sudden onset of vague and nonspecific symptoms as cough, dizziness, headache, sore throat, chills, rigors, weakness, malaise, restlessness, apprehen-

sion, myalgias, arthralgias, and fever. More than 75% of patients develop a generalized rash and, if no treatment is administered, patients may present vomiting, seizures, mental status changes, violent chills, circulatory collapse, and shock.^[1]

Concerning the laboratory examinations, leukocytosis is often seen; the chemistry profile may have an elevated anion gap metabolic acidosis and, if there is bilateral adrenal hemorrhage, hyponatremia, hyperkalemia, mild azotemia, and occasionally hypoglycemia may be found.^[21] Coagulation abnormalities are consistent with DIC and consumption of clotting factors.^[22] Thus, the clinical findings could be considered very unspecific, making a clinical diagnosis of WFS extremely challenging. Other useful information could be obtained from Gram's stains and cultures from blood, cerebrospinal fluid, urine, and sputum to have.

If clinically suspected, the CT scan of the abdomen is strongly suggested as “gold standard” in the diagnosis of adrenal hemorrhage.^[23] The therapy should be started as soon as the diagnosis is suspected, due to the lethal and fulminant course of this syndrome. It consists of antibiotic administration and, in particular, Penicillin G is the most suggested in *N meningitidis* infection, while third-generation cephalosporin can be used when the organism is unknown.^[24] The need for corticosteroidal administration was a belief fervently held by Friderichsen in his later writings.^[25] Some authors hold corticosteroid therapy as not recommended, maintaining that adrenal insufficiency is very rare and that serum cortisol levels are usually elevated in patients

Table 1**Summary of the literature reports on WFS caused by unexpected/unusual infective agents.**

Case	Pathogen	Sex	Age	Clinical course	Clinical events	Materials and methods	Outcome	Reference
1	<i>Pseudomonas aeruginosa</i>	M	2	5 d	50% body surface area second and third-degree scalding burns	Premortem MA: blood	Died	[2]
2	<i>Ewingella americana</i>	F	74	14 h	Dragging pain in her left leg	Postmortem MA: blood	Died	[11]
3	<i>Capnocytophaga canimorsus</i>	M	53	2 d	Vague abdominal pain, myalgia, and increased lethargy	Premortem MA: blood	Died	[12]
4	<i>Escherichia coli</i>	F	49	11 d	High-grade fever, nausea, urinary frequency, back pain, and rigors	Premortem MA: blood and urine	Died	[13]
5	<i>Acinetobacter calcoaceticus</i>	M	44	12 h	Severe cephalgia with stupor, petechiae on chest and arms, T: 37,8°C	Postmortem MA: brain, spinal cord, leptomeninges, kidney, lung, spleen	Died	[14]
6	DF-2 bacterium	F	57	18 h	Abdominal pain and vomiting	Postmortem MA: blood	Died	[15]
7	<i>Pasteurella multocida</i>	F	64	7.30 h	Sudden onset epigastric pain, vomiting, chills, and rigors	Premortem and Postmortem MA: blood, bile	Died	[16]
8	<i>Moraxella duplex</i>	F	63	11 d	Sore throat, general weakness, chills, fever, nausea, vomiting, semicomatose condition	Postmortem MA: meninges, CSF	Died	[17]
9	<i>Rickettsia rickettsia</i>	n.a.	n.a.	n.a.	n.a.	Postmortem IHC (ab anti- <i>Rickettsia rickettsia</i>): different tissues	Died	[18]
10								
11								
12								
13	<i>Bacillus anthracis</i>	n.a.	n.a.	n.a.	n.a.	Postmortem IHC (ab anti- <i>Bacillus anthracis</i> cell wall): different tissues		
14								
11	<i>Ehrlichia</i> sp.	M	n.a.	n.a.	n.a.	Postmortem IHC (ab anti- <i>Ehrlichia</i> spp): different tissues		
12	<i>Treponema pallidum</i>	F	fetus	n.a.	n.a.	Postmortem IHC (ab anti- <i>Treponema pallidum</i>): different tissues		
13	<i>Legionella pneumophila</i>	M	44	n.a.	n.a.	Postmortem IHC (ab anti- <i>Legionella pneumophila</i>): different tissues		
14	<i>Plesiomonas shigelloides</i>	M	43	2 d	Fever, shaking, chills, weakness, severe abdominal pain, diarrhea, vomiting	Premortem MA: blood and urine	Died	[19]
15	<i>Varicella zoster</i>	M	53	4 d	Sharp umbilical pain, general weakness, general vesicular rash	Postmortem PCR: skin lesion fluid, plasma	Alive	[20]

CSF = cerebrospinal fluid, d = days, h = hours, IHC = immunohistochemistry, MA = microbiological analysis, n.a. = not available, PCR = polymerase chain reaction.

with adrenal hemorrhage; nevertheless, documentation of dramatic clinical improvement and survival following treatment with cortisone exists.^[26] Obviously, the administration of therapy aimed to manage the clinical and laboratory tests alterations and the clinical manifestations directly related to the disease (ie, anticoagulant therapy for DIC treatment; supportive therapy in circulatory collapse and shock) is also very important.^[27]

In light of this brief overview on WFS, the presented case confirms that the collection of clinical, laboratory tests and microbiological data, together with imaging, could be useful in

practical diagnosis, even if, nowadays, the assessment of WFS is often given by postmortem findings.

The main evidence of the analyzed case was that WFS occurred in the setting of *P mirabilis* bacteremia. *P mirabilis* is described as a urease positive, highly motile bacterium, member of the Enterobacteriaceae family. It is normally found in soil, water, and sewage, and is part of the normal fecal flora. It has been related to community-acquired and nosocomial illnesses, including urinary tract infection (cystitis, pyelonephritis, and prostatitis), septicemia, and wound infections, but also in neonatal meningoencephalitis, empyema, and osteomyelitis.^[28,29]

This is the first case, reported in the literature, in which the *P mirabilis* infection was complicated by WFS. Therefore, this case report demonstrates the difficulty in diagnosing and treating the WFS and supports the theory that any bacterium which causes DIC may cause adrenal hemorrhage. Finally, the evidence of WFS related to several uncommon microorganisms should suggest to clinicians the importance to consider a potential adrenal involvement in every patient with sepsis and DIC.

Author contributions

Conceptualization: Elvira Ventura Spagnolo, Cristina Mondello, Cataldo Raffino.

Data curation: Elvira Ventura Spagnolo, Cristina Mondello, Salvatore Rocuzzo, Chiara Stassi, Luigi Cardia, Angela Grieco, Cataldo Raffino.

Formal analysis: Elvira Ventura Spagnolo, Salvatore Rocuzzo, Cataldo Raffino.

Investigation: Cataldo Raffino.

Methodology: Cristina Mondello, Luigi Cardia.

Project administration: Elvira Ventura Spagnolo.

Resources: Angela Grieco, Cataldo Raffino.

Supervision: Elvira Ventura Spagnolo, Cristina Mondello.

Validation: Elvira Ventura Spagnolo, Cristina Mondello, Cataldo Raffino.

Visualization: Chiara Stassi.

Writing – original draft: Elvira Ventura Spagnolo, Cristina Mondello.

Writing – review and editing: Luigi Cardia.

References

- Varon J, Chen K, Sternbach George L. Rupert Waterhouse and Carl Friderichsen: adrenal apoplexy. *J Emerg Med* 1998;16:643–7.
- Tormos LM, Schandl CA. The significance of adrenal hemorrhage undiagnosed Waterhouse-Friderichsen syndrome. A case series. *J Forensic Sci* 2013;58:1071–4.
- Laganà P, Dattilo G, Mondello C, et al. A case of infective endocarditis due to *Salmonella enterica* phagetype 35. First report. *Clin Ter* 2017;168:e397–400.
- Ventura Spagnolo E, Mondello C, Rocuzzo S, et al. A lethal tick-borne encephalitis (TBE) due to TBE virus in Sicily (Italy): a case of IgG+/IgM-response? *Clin Ter* 2018;169:e145–8.
- Ventura Spagnolo E, Mondello C, Cardia L, et al. Odontogenic abscess complicated by descending necrotizing mediastinitis: evidence of medical and dental malpractice. *Minerva Stomatol* 2016;65:412–5.
- Simon DR, Palese MA. Clinical update on the management of adrenal hemorrhage. *Curr Urol Rep* 2009;10:78–83.
- Levi M, Keller T, van Grop E, et al. Infection and inflammation and the coagulation system. *Cardiovasc Res* 2003;60:26–39.
- Aoyama H, Kikuchi F, Mori W. Acute, massive, haemorrhagic adrenal necrosis experimentally produced by the Shwartzman mechanism in rabbits. *Virchows Arch A Pathol Anat Histopathol* 1987;412:11–6.
- Lefebvre H, Thomas M, Duparc C, et al. Role of ACTH in the interactive/paracrine regulation of adrenal steroid secretion in physiological and pathophysiological conditions. *Front Endocrinol (Lausanne)* 2016;20:98.
- Ansurudeen I, Kopf PG, Gauthier KM, et al. Aldosterone secretagogues increase adrenal blood flow in male rats. *Endocrinology* 2014;155:127–32.
- Tsokos M. Fatal Waterhouse-Friderichsen syndrome due to *Ewingella americana* infection. *Am J Forensic Med Pathol* 2003;24:41–4.
- Cooper JD, Dorion RP, Smith JL. A rare case of Waterhouse-Friderichsen syndrome caused by *Capnocytophaga canimorsus* in an immunocompetent patient. *Infection* 2015;43:599–602.
- Khawaja J. Bilateral adrenal hemorrhage in the background of *Escherichia coli* sepsis: a case report. *J Med Case Rep* 2017;11:72.
- Ferrara SD, Bertoloni G, Terribile V. Sudden death due to waterhouse-Friderichsen syndrome and purulent Leptomeningitis caused by *Acinetobacter calcoaceticus* (*Mima polymorpha*). *Z Rechtsmed* 1977;80:73–8.
- Chaudhuri AK, Hartley RB, Maddocks AC, et al. Waterhouse-Friderichsen syndrome caused by a DF-2 bacterium in a splenectomised patient. *J Clin Pathol* 1981;34:172–3.
- Morrison U, Taylor M, Sheahan DG, et al. Waterhouse-Friderichsen syndrome complicating primary biliary sepsis due to *Pasteurella multocida* in a patient with cirrhosis. *J Clin Pathol* 1995;48:775–7.
- Lewis JF, Marshburn ET, Singletary HP, et al. Fatal meningitis due to *Moraxella duplex*: report of a case with Waterhouse-Friderichsen syndrome. *South Med J* 1968;61:539–41.
- Guarner J, Paddock CD, Bartlett J, et al. Adrenal gland hemorrhage in patients with fatal bacterial infections. *Mod Pathol* 2008;21:1113–20.
- Curti AJ, Lin JH, Szabo K. Overwhelming post-splenectomy infection with *Plesiomonas shigelloides* in a patient cured of Hodgkin's disease. A case report. *Am J Clin Pathol* 1985;83:522–4.
- Heitz AFN, Hofstee HMA, Gelinck LBS, et al. A rare case of Waterhouse-Friderichsen syndrome during primary Varicella zoster infection. *Neth J Med* 2017;75:351–3.
- Xarli VP, Steel A, Davis PJ, et al. Adrenal hemorrhage in the adult. *Medicine* 1978;57:211–21.
- Kunzer W, Schindera F, Schenk W, et al. Waterhouse-Friderichsen-syndrome. *Dtsch Med Wochenschr* 1972;97:270–3.
- Munoz Corsini L, Delgado Arnaiz C, García del Valle S, et al. Postoperative bilateral adrenal hemorrhage: correlation between clinical and radiological signs. *J Clin Anesth* 2008;20:605–8.
- Salzman MB, Rubin LG. Meningococemia. *Inf Dis Clin North Am* 1996;10:709–25.
- Friderichsen C. Waterhouse-Friderichsen syndrome. *Acta Endocrinol* 1955;18:482–92.
- Jafari HS, McCracken GH. Sepsis and septic shock: a review for clinicians. *Pediatr Inf Dis J* 1992;11:739–49.
- Carcillo JA, Davis AL, Zaritzky A. Role of early fluid resuscitation in pediatric septic shock. *JAMA* 1991;266:1242–5.
- Kim BN, Nam NJ, Kim MN, et al. Bacteraemia due to tribe proteaceae: a review of 132 cases during a decade. *Scand J Infect Dis* 2003;35:98–103.
- Chen CY, Chen YH, Lu PL, et al. *Proteus mirabilis* urinary tract infection and bacteremia: risk factors, clinical presentation, and outcomes. *J Microbiol Immunol Infect* 2012;45:228–36.