

Role of the diet as a link between oxidative stress and liver diseases

Teresa Arrigo, Salvatore Leonardi, Caterina Cuppari, Sara Manti, Angela Lanzafame, Gabriella D'Angelo, Eloisa Gitto, Lucia Marseglia, Carmelo Salpietro

Teresa Arrigo, Caterina Cuppari, Sara Manti, Gabriella D'Angelo, Eloisa Gitto, Lucia Marseglia, Carmelo Salpietro, Department of Pediatric, Gynecological, Microbiological and Biomedical Sciences, University of Messina, 98122 Messina, Italy
Salvatore Leonardi, Angela Lanzafame, Department of Medical and Pediatrics Science, University of Catania, 95123 Catania, Sicily, Italy

Author contributions: All authors of this paper have equally participated in the planning or drafting of the manuscript.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Leonardi Salvatore, MD, Department of Medical and Pediatrics Science, University of Catania, Via Santa Sofia, 64, 95123 Catania, Sicily, Italy. leonardi@unict.it

Telephone: +39-95-3782764

Fax: +39-95-378238

Received: September 4, 2014

Peer-review started: September 27, 2014

First decision: September 27, 2014

Revised: October 24, 2014

Accepted: December 1, 2014

Article in press: December 1, 2014

Published online: January 14, 2015

Abstract

Oxidative stress is caused by an imbalance between the production of reactive oxygen (free radicals) and the body's ability (antioxidant capacity) to readily detoxify the reactive intermediates or easily repair the resulting damage. An adequate diet, characterized by daily intake of foods associated with improvements

in the total antioxidant capacity of individuals and reduced incidence of diseases related to oxidation, can modulate the degree of oxidative stress. In fact, diet-derived micronutrients may be direct antioxidants, or are components of antioxidant enzymes, leading to improvement of some indicators of hepatic function. However, although their increased dietary intake might be beneficial, literature data are still controversial. This review summarizes what is known about the effects of diet nutrients on oxidative stress, inflammation and liver function. Moreover, we have analyzed: (1) the main nutritional components involved in the production and/or removal of free radicals; and (2) the role of free radicals in the pathogenesis of several hepatic diseases and related comorbidities.

Key words: Nutrition; Micronutrients; Macronutrients; Liver disease; Oxidative stress

© The Author(s) 2015. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Nutritional intake is a fundamental determinant of health. Recently, it has been observed that dietary supplementation has hepatoprotective and anti-oxidant effects. The aim of this review was to summarize the molecular changes promoted by diets and to underline the relationship between diet, oxidative stress and liver disease.

Arrigo T, Leonardi S, Cuppari C, Manti S, Lanzafame A, D'Angelo G, Gitto E, Marseglia L, Salpietro C. Role of the diet as a link between oxidative stress and liver diseases. *World J Gastroenterol* 2015; 21(2): 384-395 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v21/i2/384.htm> DOI: <http://dx.doi.org/10.3748/wjg.v21.i2.384>

INTRODUCTION

Dietary elements have long been known to play a critical role in the physiological or pathological response to tissue inflammation and oxidative stress (OS). Accordingly, deficiency or excessive production of most dietary nutrients can impair the balance between the anti- and pro-oxidant agents, also causing liver diseases. Although few studies have evaluated the histological signs in hepatic diseases after adequate and/or inadequate dietary intake^[1,2], the involvement of nutrition and OS in the pathogenesis of liver dysfunction has been assessed^[1,3,4]. In fact, specific micro- or macro- nutrients may play a critical role in liver cell integrity, influencing fibrosis, inflammation, and degree of liver steatosis caused by OS. Therefore, it is useful to understand the underlying mechanism for the production and removal of free radicals (FR) and/or other reactive species. FRs, including superoxide anions, hydroxyl radicals, and hypochlorous acid, show “two faces” in biology. They may be released in the liver as a consequence of physiological (*e.g.*, signal transduction, gene transcription, phagocytosis, hepatic detoxification, and metabolic pathways) and pathological conditions (*e.g.*, inflammation and necrosis)^[5-7]. The impairment of antioxidant defense mechanisms (*e.g.*, superoxide dismutase, catalase and glutathione peroxidase, and glucose 6-phosphate dehydrogenase) could permit enhanced free radical-induced tissue damage. Therefore, reactive oxygen species (ROS)-mediated injury to membranes, proteins, DNA and RNA, generation of pro-inflammatory cytokines, activation of spindle cells, and finally fibrogenesis can occur^[8,9]. The main targets of OS are the endoplasmic reticulum (ER) and mitochondria.

The ER regulates the synthesis and release of membrane proteins. The maintenance of its function requires high concentrations of intra-ER Ca^{2+} . Several injuries induce a decrease in physiologically high intra-ER Ca^{2+} levels that result in impaired ER function, also known as “ER stress”, promoting apoptosis, hepatic stellate and Kupffer cells recruitment, and synthesis of inflammatory cytokine-inducible factors^[10].

Although mitochondria can control the oxidative balance, under continuous stressful stimuli they collapse and become producers of oxidative damage^[11]. In fact, mitochondrial fatty acids are normally transferred to the β -oxidation pathway. Fatty acid overload can lead to an imbalance between increased delivery of electrons to the respiratory chain and their decreased outflow from this chain, causing accumulation of ROS and peroxidation products. Thus, mitochondria represent the main source of ROS. Furthermore, ROS attack mitochondrial DNA and promote accumulation of DNA mutations, which, in turn, lead to further ROS synthesis^[11]. Additionally, by inducing collagen gene expression, lipid peroxidation products can stimulate fibrogenesis and apoptosis. The released apoptotic bodies can activate hepatic stellate cells and convert them, through transforming growth factor β 1, platelet-derived growth factor and endothelial growth factor, into myofibroblasts. Moreover,

apoptotic bodies can also stimulate Kupffer cells to generate ROS, which further enhance apoptosis, and the release of cytokines and chemokines that contribute to the activation of hepatic stellate cells into myofibroblasts^[12]. Myofibroblasts, through nicotinamide adenine dinucleotide phosphate, can also induce the ROS synthesis, and the release of adhesion molecules for T lymphocytes and natural killer T-cells. Finally, myofibroblasts generate an extracellular matrix resistant to degradation mediated by metalloproteinases^[13].

The consequence of these numerous interactive cellular and molecular pathways is to perpetuate and to enhance hepatic tissue injury. Specific dietary nutrients can interfere in the aforementioned pathways, potentially modulating the conversion of highly reactive FR to relatively inert radicals.

NUTRITION AND HEPATIC OXIDATIVE STRESS

Literature data support the usefulness of anti-oxidant activities of micro- and macro- nutrients to prevent many human diseases (*e.g.*, neoplasia, inflammation, autoimmune diseases). On the other hand, growing evidence suggests that hypo- and hyper-nutrition are associated with a higher likelihood of cellular oxidation^[14], mediated by specific nutrient pathways.

Here, we discuss the role of micro- and macro- nutrients intake on the development hepatic disease.

Proteins

Proteins, including amino acids (*e.g.*, arginine, citrulline, glycine, histidine, and taurine), small peptides (carnosine), and nitrogenous metabolites (*e.g.*, creatine and uric acid) directly scavenge ROS and also inhibit inducible-nitric oxide synthase (iNOS) expression in various cell types, including hepatocytes^[15]. However, when the capacity of this antioxidant system decreases, the level of inactivated ROS rises. ROS-mediated modification might alter both protein structure and function. OS may induce reversible and irreversible changes in proteins. Reversible alteration, generally involving cysteine, can modulate the function of a protein. Irreversible modification, usually lysine, results in a permanent loss of function and may contribute to the degradation and the accumulation of proteins into cytoplasmic inclusions^[16]. Moreover, oxidized proteins are highly susceptible to proteolytic attack by proteasomes. Thus, a dietary deficiency of protein is associated with decreased synthesis of antioxidant enzymes and an increased superoxide anion release. Likewise, protein malnutrition can also cause steatosis, modulating the expression of lipolysis and lipid utilization genes in liver^[17]. Proteins are involved in several mechanisms of hepatic fat storage. In fact, an adequate protein diet was associated with: (1) major metabolic rates and mitochondrial oxygen consumption; (2) increased β -oxidation of fatty acids; and (3) elevated bile acids levels which, in turn, inhibit lipogenesis^[18].

Finally, proteins in the hepatic tissue, produce a large amount of energy.

In animal models, it has been reported that insufficient protein intake causes a deficiency of micro-nutrients (zinc) and serum albumin levels^[14,19]. In fact, plasma free iron values are elevated in malnourished patients. Some studies suggested that the increased tissue free iron concentrations likely result from low concentrations of hepatic iron-binding proteins (ferritin, transferrin, and lactoferrin)^[20].

An increment in dietary protein content has been noted to prevent the likelihood of liver fat accumulation in excessive fat intake in humans^[21].

However, recently, an increased protein intake has been shown to stimulate the generation of ROS, and lipid peroxidation in human polymorphonuclear leukocytes and mononuclear cells^[22].

For these reasons, a high-protein and low-carbohydrate diet is not typically used in children because of their low energy intake, which might compromise the child's growth^[23]. However, this type of diet is preferred, for a short time, to induce a rapid weight loss in obese children affected by non-alcoholic fatty liver disease (NAFLD)^[23].

In conclusion, although the exact mechanism remains to be elucidated, an adequate quality and quantity of protein intake might be helpful in fatty liver disease.

Carbohydrates

Epidemiologic studies have implicated foods containing high concentrations of carbohydrates in the etiology of liver diseases. In fact, high carbohydrate intake, by activation of specific transcription factors, promotes hepatic steatosis and insulin resistance. The most common simple carbohydrates, glucose and fructose, can result in relatively high glycemic index values. High glycemic index foods stimulate excessive and prolonged insulin secretion, which increases deposition of fats and leads to major serum non-esterified free fatty acid values into circulation and in the hepatocytes mass. According to these findings, subjects affected by hepatic disease should avoid high carbohydrate intake, especially fructose^[24], which, by inducing depletion of adenosine triphosphate (ATP), causes arrest in protein production, favors inflammatory proteins release, alters endothelial function, and stimulate OS^[25]. Several adult studies have found links between higher consumption of total carbohydrates and nonalcoholic steatohepatitis (NASH) and the metabolic syndrome. Although fructose's role in human NAFLD is unknown, it is hypothesized that in addition to OS, lipid peroxidation, cytokine activation, nitric oxide (NO) and ROS, endogenous toxins of fructose metabolites can further lead to hepatic fat accumulation^[26]. Fructose promotes intestinal permeability leading to portal overload of endotoxins, pro-inflammatory factors [tumor necrosis factor- α : TNF- α], and fatty acids in the liver^[27].

Lower serum high-density lipoprotein levels and

higher triglyceride values were previously associated with major fructose intake in children affected by NAFLD^[27]. Moreover, mineral deficiency, metabolic imbalance, and higher release of ROS can further favor fructose-induced NAFLD^[27]. Recently, a study in normal-weight and overweight children showed that total fructose intake was the only dietary factor that significantly predicted low-density lipoprotein particle size^[28].

However, it has been also reported that changes in the quality of carbohydrates intake did not influence hepatic functionality. In fact, isocaloric exchange of fructose for other carbohydrates seems not induce NAFLD changes in healthy subjects^[29].

Lipids

Fat is an important component of the normal human diet. It is a source of energy and provides essential fatty acids and fat-soluble vitamins. However, saturated fatty acids (SFAs) and polyunsaturated fatty acids (PUFAs) can have adverse effects on human health. Although SFAs promote OS-resistant states in the liver and protect it against OS^[30], studies have demonstrated that SFAs are more toxic than unsaturated fats. High SFA intake leads to a proinflammatory status resulting from an imbalance in lipid signaling pathways and increased production of inflammatory cytokines (*e.g.*, TNF- α , IL-6)^[31]. Moreover, SFAs promote endoplasmic reticulum stress as well as hepatocyte injury. In fact, accumulation of SFAs in the liver leads to impaired mitochondrial metabolism and increased markers (ROS) related to endoplasmic reticulum stress. This latter, in turn, contributes to hepatic apoptosis progression^[32].

The accumulation of SFAs in the hepatocytes can promote apoptosis by intrinsic and extrinsic pathways. The intrinsic pathway of cell death includes: (1) ROS-induced stress that affects the endoplasmic reticulum, mitochondrial membranes, and lysosomes; and (2) lipid peroxidation that increases the serum ROS levels^[33]. The extrinsic pathway is mediated by death ligands, such as Fas (a key death receptor belonging to the TNF-receptor family) and TRAIL (TNF-related apoptosis-inducing ligand), which subsequently stimulate TNF- α production. TNF- α consequently induces the upregulation of pro-apoptotic molecules and, finally, cell death^[34].

If overeating SFA promotes hepatic damage and visceral fat storage, an excess energy from PUFA may contribute to perpetuate liver injury. PUFAs are essential fatty acids, which are crucial for normal growth and health, and are not synthesized in the body of mammals. PUFAs exhibit anti-inflammatory action by suppressing pro-inflammatory cytokine production, macrophages and hepatocytes^[35]. ω -6 PUFAs (*e.g.*, linoleic acid, alpha-linoleic acid, and arachidonic acid) contribute to the regulation of fatty acid synthesis and oxidation in the liver. However, high intake of ω -6 PUFAs may increase lipid peroxidation, iNOS expression, FR production and oxidative DNA damage in many cell types such

as macrophages, muscle and liver cells. Higher fatty acids levels in portal or systemic circulation promote visceral and hepatic fat deposits, mediated by decreased synthesis of lipoproteins and export of lipids from the liver. Subsequently, liver fatty infiltration may also cause decreased fatty acid oxidation and hepatic steatosis^[36]. ω -3 PUFAs (*e.g.*, α -linolenic acid, eicosapentaenoic acid, docosahexaenoic acid) inhibit lipogenesis and stimulate fatty acid oxidation in the liver^[23]. In rodents, ω -3 PUFAs may influence body composition and obesity. Major intake of ω -3 PUFAs may decrease fat accumulation, mainly visceral fat, and reduce body weight when already obese^[37]. It has been suggested that ω -3 PUFAs may activate a metabolic change in adipocytes, including increased β -oxidation, suppressed lipogenesis, decrease fat accumulation and higher apoptosis. Additionally, ω -3 PUFAs seem to reduce prostaglandin synthesis, even more than ω -6 PUFAs^[38].

Liver-specific responses, such as regulation of blood glucose homeostasis, sinusoidal blood flow within the liver, properties of the trans-endothelial barrier within the liver, synthesis and release of important other mediators like cytokines, growth factors or NO, and liver fibrogenesis, are mediated or regulated by prostaglandin E2 (PGE2)^[39]. Appropriate PGE2 levels are generated through the activation of constitutive cyclooxygenase-1 (COX-1) in hepatocytes. PGE2 plays a crucial role in liver pathophysiology *via* essentially hepatoprotective functions, such as inhibiting the generation of ROS, preventing leukocyte migration, improving hepatic insulin and lipid metabolism and regulating the production of inflammatory cytokines^[40]. On the other hand, PGE2 also induces the expression of inflammatory cytokines, which can, in turn, enhance the production of ROS^[41].

Previously, Mater *et al.*^[42] suggested that ω -3 PUFAs combines with the specific site on the COX-1 enzyme that converts ω -6 PUFAs into prostaglandins. Moreover, they can also act as the precursor of prostaglandins; however, their activity is 2-50 times lower than the prostaglandin produced by ω -6 PUFAs. It has also been observed that ω -3 PUFAs influence the synthesis of inflammation resolution mediators by neutrophils^[43]. In animals, the anti-inflammatory effect was correlated with overexpression of antioxidant genes (glutathione-S-transferases, uncoupling protein-2 and Mn-SOD)^[44]. In humans, however, data are controversial. An insufficient ω -3 PUFAs intake was promotes susceptibility to hepatic inflammation in children with NAFLD^[45]. Although it has been reported that ω -3 PUFAs ameliorate hepatic circulation, mediated by suppression of local and systemic proteins, including high-mobility group box 1 (HMGB1)^[46-48], an adequate dietary intake of ω -3 PUFA did not reflect systemic pro-inflammatory cytokines and protein levels^[49].

It has been reported that diet integration with ω -3 PUFAs might facilitate hepatic metabolic adaptation from in utero nutrition to the postnatal diet, by increasing fatty

acid oxidation and modifying glucose and amino acids to anabolic pathways^[50]. In addition to normocaloric/normolipidic diet, ω -3 PUFAs treatment correlated with the best metabolic parameters results (lower rise of serum triglycerides, glycemia, and cholesterol levels in serum) and reversed the liver histopathological results^[51]. Recently, Chahal *et al.*^[37] assessed that ω -3 PUFAs supplementation was not significantly effective in treating hypertriglyceridemia in pediatric patients. Although the mechanism remains unclear, it has been proposed that ω -3 PUFAs, in addition to total parenteral nutrition (TPN) and weight reduction therapy^[52], reverse or improve abnormal liver tests in children^[53]. To date, studies are insufficient regarding the types, amount, and duration of the intake of PUFAs in children^[54].

Vitamins

Many vitamins play an important role in preventing radical induced cytotoxicity by inhibition of iNOS activity, its gene transcription and NO production. Vitamins also directly scavenge ROS and upregulate the activities of antioxidant enzymes^[15].

Among them, vitamin E (α -tocopherol) is thought to be one of the most important micronutrients that inhibit ROS-induced release of lipid peroxy radicals, protecting cells from pro-oxidants and OS. Vitamin E, acting as a chain-breaking antioxidant, prevents the propagation of FR in membranes and in plasma lipoproteins. When peroxy radicals are formed, these quickly react with vitamin E (Vit EOH). The hydroxyl group of tocopherol reacts with the peroxy radical to form the corresponding lipid hydroperoxide and tocopheryl radical (Vit E-O). This latter, by binding vitamin C, returns vitamin E to its reduced state^[55]. The interaction of vitamins E and C has led to the hypothesis of the "antioxidant network", also known as "vitamin E recycling", by which the antioxidant function of oxidized vitamin E is continuously restored by other antioxidants^[56]. During phlogosis, this mechanism also reduces mast cell activation. Mast cells are activated by oxidized lipoproteins, resulting in increased expression of inflammatory cytokines and suggesting the reduction of oxidation of low-density lipoprotein by vitamin E^[57]. Therefore, a dietary deficiency of vitamin E reduces the activities of antioxidant enzymes (such as liver GSH peroxidases, glutathione reductase, and catalase) and leads to increased hepatic lipid peroxidation. Fortunately, all these negative effects can be reversed by dietary vitamin E supplementation^[58]. However, the efficacy of vitamin E remains controversial in the treatment of liver diseases, including NAFLD. Several experimental studies assessed the potential protective role of vitamin E and showed that it improves the clinical symptoms of NAFLD^[59], enhances glucose metabolism^[60], and correlates with lower serum transaminase levels^[61]. Moreover, Vitamin E, decreases histopathological damage (including degree of steatosis, inflammation and fibrosis) in adults and

children^[62-64]. However, other authors did not confirm these data^[65-67].

Currently, randomized controlled trials in children have not demonstrated uniformly the beneficial effects of vitamin E on the long-term outcome of NAFLD patients. The TONIC trial did not find a significant correlation between the vitamin E and control groups in improving serum alanine aminotransferase levels, steatosis or hepatic inflammation. However, increased resolution of hepatocyte ballooning was observed in the vitamin E subjects^[64].

In humans, vitamin C, a water-soluble electron donor, plays a protective role against FR-induced OS. Vitamin C inhibits peroxidation of membrane phospholipids and acts as a scavenger of FR (superoxide, singlet oxygen, and hydroxyl radicals) and is also required for one-electron reduction of lipid hydroperoxyl radicals *via* the vitamin E redox cycle^[67]. Vitamin C prevents hepatic storage of 8-hydroxydeoxyguanosine, a marker of DNA injury, and hinders hepatocellular growth. Moreover, vitamin C arrests bacteria internalization and translocation by increasing the transepithelial membrane resistance^[68] and enhancing the ability of neutrophils to kill bacteria. In the presence of bacteria, ascorbate levels in neutrophils increase to protect these cells against damage by ROS that they previously produced^[69].

In an experimental model, ascorbate's anti-fibrotic action, attributed to decrease in the oxidative stress, hepatic stellate cells activation, cytotoxicity and mRNA expression of fibrotic genes, has been also been reported^[70]. The aforementioned positive effects of vitamins could provide a rationale for dietary vitamin C intake in individuals. However, the supplementation of vitamin C as further liver treatment is still controversial. A Cochrane database meta-analysis, analyzing six trials that used a combination of selenium, vitamin C and vitamin E to evaluate their effects on the NAFLD, found no evidence to support or refute them as useful treatments^[71]. Intervention studies with vitamin C have shown no change in markers of oxidation or in clinical benefit. In fact, in children, authors reported no additional effects of using vitamin C or E on weight loss and a possible histological improvement in patients affected by NAFLD^[1,72]. Perhaps these controversial data are related to different administration doses. Dose concentration studies of vitamin C in healthy people showed a sigmoidal relationship between oral dose and plasma and tissue vitamin C concentrations. Hence, optimal dosing is critical for intervention studies using vitamin C^[73].

Vitamin B₁₂ is an essential cofactor that plays important roles in one-carbon metabolism, which is required for the maintenance of intracellular DNA synthesis and methylation. In fact, vitamin B₁₂, serving as a cofactor for methionine synthase, cystathionine synthase, and cystathionase, and as a substrate (5-methyl-tetrahydrofolate) for methionine synthase, helps to reduce

the risk of OS-mediated homocysteine^[74]. The liver is the principal storage site of vitamin B₁₂. Consequently, serum Vitamin B₁₂ levels reflect liver function. In fact, it has been reported that vitamin B₁₂ and hepatic enzyme serum levels are correlated, especially in alcohol-dependent liver disease^[74].

Vitamin B₁₂, through epigenomic mechanisms related to imbalanced acetylation/methylation, influences cell proliferation, differentiation and apoptosis. In addition, vitamin B₁₂, by ER stress stimulation, impairs fatty acid oxidation and energy metabolism in the liver^[75]. Therefore, increased serum vitamin B₁₂ levels have been attributed to the release of the vitamin from the liver during hepatic necrosis and decreased hepatic synthesis of transcobalamins II, an essential element for tissue binding of vitamin B₁₂^[75].

Minerals

Studies on most micronutrients are complicated by the fact that the nutrients have several roles.

Selenium, a cofactor of numerous enzymes (*e.g.*, glutathione peroxidase, selenoprotein P, and other selenoproteins) has a protective role against peroxidative and/or FRs damage and mitochondrial dysfunction^[76]. In humans, deficiency of selenoproteins causes liver necrosis and hepatic cell death by OS. Furthermore, low serum selenium levels have been found in patients with chronic liver disease^[77].

Magnesium is involved in ATP-mediated reactions. Deficiency of dietary magnesium reduces glutathione reductase activity, and results in generation of ROS and increased susceptibility to lipid peroxidation, and marked lesions in tissues (*e.g.*, skeletal muscle, brain, and kidney)^[78].

Manganese is a component of several enzymes involved in fatty acid and cholesterol biosynthesis, as well as mitochondrial pathways. Manganese is a cofactor for a number of enzymes important for intermediary metabolism, including the hepatic urea cycle enzyme arginase. There are few well-described cases of manganese deficiency in the medical literature^[14] and how manganese affects normal metabolism in the liver remains unclear^[79].

Copper, zinc and manganese are indispensable metals for the activities of Cu, Zn-SOD and Mn-SOD, respectively^[80].

The liver plays an important role in the disposition of copper, which can be used for protein and energy production. Consequently, abnormal copper metabolism can also cause oxidative damage and hepatotoxicity. Mitochondria are the first responders involved in copper homeostasis. In an animal model, Cu²⁺ induced a concentration and time-dependent rise in mitochondrial ROS formation, lipid peroxidation, cytochrome c expulsion, mitochondrial swelling and collapse, and finally cell death signaling. Interacting with respiratory complexes (I, II, and IV), Cu²⁺ caused decreases the ATP concentration

and the ATP/ADP ratio mitochondria, favoring liver toxicity^[81]. Synthesis of ROS, tissue failure and hepatocyte death are also observed in human disease progression, such as in Wilson disease (WD). Liver damage in WD appears to involve two different pro-oxidants mechanisms: synthesis of ROS, mediated by the Haber-Weiss reaction^[82], and apoptosis through the activation of acid sphingomyelinase and consequent production of ceramide^[83,84].

The liver plays a central role in zinc homeostasis, removing it from albumin in the blood and distributing it to the body as needed. Zinc acts as an essential cofactor for enzymes that are necessary to counteract hepatic OS^[85]. Cirrhotics show increased urinary zinc loss and can become zinc deficient^[77].

Liver is an important site for iron, a cofactor for important biological biochemical reactions, including the transport of oxygen and electrons *via* cells, oxidative phosphorylation, energy production, DNA synthesis, cell growth or apoptosis and gene expression^[20]. However, iron's bioavailability is limited because its accumulation can enhance OS and related toxic effects. Therefore, iron deficiency and overload are generally regarded as causes of diseases involving OS and lipid peroxidation^[86]. Patients with iron deficiency anemia are more sensitive to agents that induce OS^[87,88]. On the other hand, daily iron supplementation resulted in increased lipid peroxidation and abnormal iron accumulation; although intermittent supplementation (once every 3 d) alleviated these effects^[89]. It is thought that iron overload, mediated by mobilization of peripheral fat to the liver and development of hyperinsulinemia, promotes insulin resistance. Therefore, the syndrome of "insulin resistance-associated iron overload" was hypothesized in the presence of unexplained hepatic iron overload in patients with insulin resistance^[90].

Increased iron storage has been linked with more advanced stages of NAFLD, *via* increased inflammation and oxidative stress^[91].

Increased iron accumulation in NAFLD might be the results of decreased serum iron export protein levels, ferroportin, induced by the decrease of the iron regulatory peptide, and hepcidin. Generally, synthesis of hepcidin is upregulated by phlogosis and increased iron stores. Hepcidin in turn leads to the degradation of ferroportin and iron release^[92]. In patients with NAFLD, increased expression of hepcidin and lower expression of ferroportin could favor iron accumulation and, consequently, OS and inflammation^[93]. Iron overload is also very common in many types of non-biliary cirrhosis, and in end stage liver disease, including hemochromatosis. When liver iron overload is excessive, OS, involving production of iron catalyzed oxygen radicals, represents the main mechanism of liver injury, DNA alterations, and higher risk of cancer in patients affected by hemochromatosis^[94].

Although the degree of necro-inflammation did

not differ, iron stores were reported to be also elevated in patients with chronic hepatitis C compared to those with chronic hepatitis B^[95]. HCV infection is probably associated with increased hepatic iron concentration and deposition (Kupffer cells and portal macrophages), higher serum transferrin saturation, and serum ferritin levels. Additionally, serum iron values were reported to be correlated with progression of liver disease and degree of hepatocyte necrosis^[96]. Moreover, iron overload can activate hepatic stellate cells and promote the synthesis of collagen, contributing to hepatic fibrogenesis^[97].

Probiotics

Probiotics, also known as "good bacteria" or "helpful bacteria," are live microorganisms (*e.g.*, bacteria), which are beneficial to health and are either the same as or similar to microorganisms found naturally in the human body (*Lactobacillus* or *Bifidobacterium*)^[98].

Probiotics play a pivotal role in NAFLD and NASH, obesity-related hepatocarcinogenesis, alcohol-related disorders, portal hypertension and obstructive jaundice^[6,99,100].

Patients with hepatic diseases showed an impaired gut-liver axis, which contributes to increase blood levels of endotoxemia (lipopolysaccharides, LPS) and chronic low-grade inflammation. Generally, LPS disseminates into the systemic circulation in two different ways: *via* a portal vein or through the lymphatic system. Underlying liver disease caused patients to report an increased gut permeability which promotes bacterial overgrowth and translocation, and increased expression of pro-inflammatory molecules (iNOS, ROS), which in turn promote major gut and sinusoidal permeability, increased pro-inflammatory cytokines and cells (*e.g.*, neutrophil), and mitochondrial damage^[7,99,101].

The use of probiotics to manage liver injury is attributed to a variety of their health benefits. Probiotics, which increase cellular permeability and compete with pathogens for binding, improve colonization resistance to gut pathogens by reinforcing the mucosal barrier and restoring normal gut micro-ecology^[102,103]. Probiotics can activate and modulate the immune system, and reinforce gut defense by immune exclusion, elimination and regulation^[104,105]. The link between gut microbiota, liver inflammation, and immune system involves toll-like receptors (TLRs), which are important mediators between the environment and the immunological response^[106] and endogenous substances, such as short-chain fatty acids and HMGB1. TLRs involved in the pathogenesis of NASH are TLR2 (for lipoproteins and glycolipids in bacteria adhering to myeloid dendritic cells, mast cells or monocytes), TLR4 (for palmitic- stearic and lauric-acid, and LPS of B cells myeloid dendritic cells, mast cells, monocytes and intestinal epithelium) and TLR9 (for unmethylated CpG DNA-bacterial particles)^[100,107].

Additionally, probiotics reduce hepatic triglyceride contents, ameliorate adipose tissue inflammation^[108],

and induce anti-oxidative enzymes that prevent the progression of NASH to hepatocellular carcinoma^[58]. In animal models with portal hypertension, by reducing bacterial translocation, probiotics decreased the OS and/or increase vasodilator factors leading to improved endothelial dysfunction in the mesenteric artery^[109]. Moreover, an experimental study investigated the role of probiotics in obstructive jaundice. The authors reported that probiotics (*Lactobacillus plantarum*), by activating the protein kinase C pathway, could decrease intestinal epithelial cell apoptosis, reduce OS, and prevent tight junction disruption in biliary obstruction^[101].

Although, Cochrane meta-analysis^[110] did not approve or refute the use of probiotics as a therapeutic option for patients with NAFLD/NASH, several recent studies showed encouraging preliminary results^[111,112]. Compared with controls, patients with liver diseases had initially decreased *Bifidobacterium* and *Lactobacillus* levels and their serum ALT, AST and GGT values were elevated significantly. After treatment, the group who received probiotics had significantly increased *Bifidobacterium* and *Lactobacillus* levels and decreased serum liver enzyme levels compared with patients receiving the placebo^[113]. This evidence confirmed both gut-liver axis malfunction and the possible useful role of probiotics in the treatment of liver diseases.

TPN AND HEPATIC OXIDATIVE STRESS

TPN is life saving in patients with clinical problems that preclude enteral diet for a long period. However, long-term TPN-related complications, especially liver dysfunction, have been reported and confirmed by a biological significant increase of cytolysis^[114,115]. Several possible mechanisms of TPN-induced liver dysfunction have been hypothesized. Firstly, TPN promotes alteration of some trace elements in hepatocytes. Depletion in the hepatic copper concentration might cause a decrease in the activity of antioxidant and detoxifying enzymes. A parallel depletion of zinc, a cofactor of tissue matrix metalloproteinase that degrades collagen, could induce the accumulation of the extracellular matrix, finally leading to the development of hepatic fibrosis and cirrhosis^[116,117].

Secondly, TPN administration favors hepatic lipid accumulation, especially in children^[118]. In addition to increased lipid synthesis, accelerated mobilization of fat deposits, impaired fatty acids oxidation and lipid accumulation leads to steatosis, mitochondrial and ER damage, activation of caspases, and consequent Fas ligand/TNF- α -mediated apoptosis. TPN also causes the depletion of carnitine, a compound necessary for the transfer of free fatty acids from the hepatic cytoplasm into the mitochondria. The cytosolic concentration of choline, a nutrient for lipoprotein release, also decreases, promoting lipid storage in hepatic cells^[119,120]. Moreover, TPN favors lipid peroxidation by providing PUFAs and

perpetuating lipid peroxidation^[121].

Thirdly, TPN might lead to impairment of the gastrointestinal immune system and the development of infections. The factors facilitating this process include: (1) abnormal proliferation and translocation of bacteria^[122]; (2) decreased neutrophilic opsonic activity^[123]; (3) atrophy of intestinal mucosa and related reduction of serum IgA levels; and (4) reduction of T-helper and IL-2 producing cells^[124].

Finally, light exposure to nutrient mixtures affects hepatobiliary responses and histological changes. When TPN was administered intravenously, no damage was noted. Additionally, the severity of TPN-liver damage could be correlated with the duration of TPN administration^[125].

In contrast to these data, an observational study assessed that, although the TPN-group showed some signs of increased OS, there were no signs for oxidative damage, compared with control-group. Moreover, the activity of the underlying disease was not correlated to increased OS^[126]. Other authors confirmed that an increase in OS bio-markers are not necessarily related to the route of pharmaconutrition (TPN); however, it might occur independently^[127].

In the light of these data, to reduce the risk of liver disease related to TPN, several management strategies have been proposed recently, including reformulation of standardized parental nutrition. It has also been proposed that an optimal dose and type of parental lipid should be provided to minimize hepatic injury^[128].

CONCLUSION

Although it is very difficult to correlate the biochemistry of a dietary intake with the pathophysiology of hepatic disease, diet significantly attenuates the relationship between OS and liver inflammation. In fact, dietary interventions may reduce the impact of hepatic diseases and could be useful in the treatment and prevention of progression to more severe disease. However, although the anti-oxidant properties of the diet are known, investigations into the relationship between the diet and OS are still limited, and most studies were conducted on a single diet element. We believe that the evaluation of combined and parallel roles of nutrients on inflammation and OS might be more helpful. Additionally, the potential correlation between diet nutrients and serological, histopathological, and molecular markers should be investigated.

Finally, the knowledge of enzymatic and non-enzymatic oxidative defense mechanisms will serve as a guiding principle for establishing the most effective nutrition intake to ensure adequate biological support, especially in patients affected by liver diseases. Investigations into the relationship between the dietary intake and OS mediators, which are enhance the inflammatory response directly and/or indirectly, are needed.

REFERENCES

- 1 **Nobili V**, Manco M, Devito R, Di Ciommo V, Comparcola D, Sartorelli MR, Piemonte F, Marcellini M, Angulo P. Lifestyle intervention and antioxidant therapy in children with nonalcoholic fatty liver disease: a randomized, controlled trial. *Hepatology* 2008; **48**: 119-128 [PMID: 18537181 DOI: 10.1002/hep.22336]
- 2 **Huang MA**, Greenson JK, Chao C, Anderson L, Peterman D, Jacobson J, Emick D, Lok AS, Conjeevaram HS. One-year intense nutritional counseling results in histological improvement in patients with non-alcoholic steatohepatitis: a pilot study. *Am J Gastroenterol* 2005; **100**: 1072-1081 [PMID: 15842581]
- 3 **Cesaratto L**, Vascotto C, Calligaris S, Tell G. The importance of redox state in liver damage. *Ann Hepatol* 2004; **3**: 86-92 [PMID: 15505592]
- 4 **Jablonowska E**, Tchórzewski H, Lewkowicz P, Kuydowicz J. Reactive oxygen intermediates and serum antioxidative system in patients with chronic C hepatitis treated with IFN- α and thymus factor X. *Arch Immunol Ther Exp (Warsz)* 2005; **53**: 529-533 [PMID: 16407785]
- 5 **Dalgıç B**, Sönmez N, Biberoğlu G, Hasanoğlu A, Erbaş D. Evaluation of oxidant stress in Wilson's disease and non-Wilsonian chronic liver disease in childhood. *Turk J Gastroenterol* 2005; **16**: 7-11 [PMID: 16252181]
- 6 **Manti S**, Marseglia L, D'Angelo G, Filippelli M, Cuppari C, Gitto E, Romano C, Arrigo T, Salpietro C. Portal hypertension as immune mediate disease. *Hepat Mon* 2014; **14**: e18625 [PMID: 24976841 DOI: 10.5812/hepatmon.18625]
- 7 **Manti S**, Romano C, Chirico V, Filippelli M, Cuppari C, Loddo I, Salpietro C, Arrigo T. Nonalcoholic Fatty liver disease/non-alcoholic steatohepatitis in childhood: endocrine-metabolic "mal-programming". *Hepat Mon* 2014; **14**: e17641 [PMID: 24829591 DOI: 10.5812/hepatmon.17641]
- 8 **Pessayre D**, Berson A, Fromenty B, Mansouri A. Mitochondria in steatohepatitis. *Semin Liver Dis* 2001; **21**: 57-69 [PMID: 11296697]
- 9 **Younossi ZM**, Diehl AM, Ong JP. Nonalcoholic fatty liver disease: an agenda for clinical research. *Hepatology* 2002; **35**: 746-752 [PMID: 11915019]
- 10 **Fu S**, Yang L, Li P, Hofmann O, Dicker L, Hide W, Lin X, Watkins SM, Ivanov AR, Hotamisligil GS. Aberrant lipid metabolism disrupts calcium homeostasis causing liver endoplasmic reticulum stress in obesity. *Nature* 2011; **473**: 528-531 [PMID: 21532591 DOI: 10.1038/nature09968]
- 11 **Murphy MP**. How mitochondria produce reactive oxygen species. *Biochem J* 2009; **417**: 1-13 [PMID: 19061483 DOI: 10.1042/BJ20081386]
- 12 **Bhagal RH**, Curbishley SM, Weston CJ, Adams DH, Afford SC. Reactive oxygen species mediate human hepatocyte injury during hypoxia/reoxygenation. *Liver Transpl* 2010; **16**: 1303-1313 [PMID: 21031546 DOI: 10.1002/lt.22157]
- 13 **Yoshiji H**, Kuriyama S, Miyamoto Y, Thorgeirsson UP, Gomez DE, Kawata M, Yoshii J, Ikenaka Y, Noguchi R, Tsujinoue H, Nakatani T, Thorgeirsson SS, Fukui H. Tissue inhibitor of metalloproteinases-1 promotes liver fibrosis development in a transgenic mouse model. *Hepatology* 2000; **32**: 1248-1254 [PMID: 11093731]
- 14 **Fang YZ**. Free radicals and nutrition. Theory and application of free radical biology. Beijing: Scientific Press, 2002: 647
- 15 **Wu G**, Meininger CJ. Regulation of nitric oxide synthesis by dietary factors. *Annu Rev Nutr* 2002; **22**: 61-86 [PMID: 12055338]
- 16 **Lu SC**. Antioxidants in the treatment of chronic liver diseases: why is the efficacy evidence so weak in humans? *Hepatology* 2008; **48**: 1359-1361 [PMID: 18697215 DOI: 10.1002/hep.22463]
- 17 **Uebanso T**, Taketani Y, Fukaya M, Sato K, Takei Y, Sato T, Sawada N, Amo K, Harada N, Arai H, Yamamoto H, Takeda E. Hypocaloric high-protein diet improves fatty liver and hypertriglyceridemia in sucrose-fed obese rats via two pathways. *Am J Physiol Endocrinol Metab* 2009; **297**: E76-E84 [PMID: 19435858 DOI: 10.1152/ajpendo.00014.2009]
- 18 **Duran-Sandoval D**, Cariou B, Percevault F, Hennuyer N, Grefhorst A, van Dijk TH, Gonzalez FJ, Fruchart JC, Kuipers F, Staels B. The farnesoid X receptor modulates hepatic carbohydrate metabolism during the fasting-refeeding transition. *J Biol Chem* 2005; **280**: 29971-29979 [PMID: 15899888]
- 19 **Rassaf T**, Preik M, Kleinbongard P, Lauer T, Heiss C, Strauer BE, Feelisch M, Kelm M. Evidence for in vivo transport of bioactive nitric oxide in human plasma. *J Clin Invest* 2002; **109**: 1241-1248 [PMID: 11994413]
- 20 **Wang J**, Pantopoulos K. Regulation of cellular iron metabolism. *Biochem J* 2011; **434**: 365-381 [PMID: 21348856 DOI: 10.1042/BJ20101825]
- 21 **Bortolotti M**, Kreis R, Debard C, Cariou B, Faeh D, Chetiveaux M, Ith M, Vermathen P, Stefanoni N, Lé KA, Schneiter P, Krempf M, Vidal H, Boesch C, Tappy L. High protein intake reduces intrahepatocellular lipid deposition in humans. *Am J Clin Nutr* 2009; **90**: 1002-1010 [PMID: 19710199 DOI: 10.3945/ajcn.2008.27296]
- 22 **Mohanty P**, Ghanim H, Hamouda W, Aljada A, Garg R, Dandona P. Both lipid and protein intakes stimulate increased generation of reactive oxygen species by polymorphonuclear leukocytes and mononuclear cells. *Am J Clin Nutr* 2002; **75**: 767-772 [PMID: 11916766]
- 23 **Spear BA**, Barlow SE, Ervin C, Ludwig DS, Saelens BE, Schetzina KE, Taveras EM. Recommendations for treatment of child and adolescent overweight and obesity. *Pediatrics* 2007; **120 Suppl 4**: S254-S288 [PMID: 18055654]
- 24 **Agius L**. High-carbohydrate diets induce hepatic insulin resistance to protect the liver from substrate overload. *Biochem Pharmacol* 2013; **85**: 306-312 [PMID: 23022226 DOI: 10.1016/j.bcp.2012.09.019]
- 25 **Lanaspa MA**, Sanchez-Lozada LG, Cicerchi C, Li N, Roncal-Jimenez CA, Ishimoto T, Le M, Garcia GE, Thomas JB, Rivard CJ, Andres-Hernando A, Hunter B, Schreiner G, Rodriguez-Iturbe B, Sautin YY, Johnson RJ. Uric acid stimulates fructokinase and accelerates fructose metabolism in the development of fatty liver. *PLoS One* 2012; **7**: e47948 [PMID: 23112875 DOI: 10.1371/journal.pone.0047948]
- 26 **Nomura K**, Yamanouchi T. The role of fructose-enriched diets in mechanisms of nonalcoholic fatty liver disease. *J Nutr Biochem* 2012; **23**: 203-208 [PMID: 22129639 DOI: 10.1016/j.jnutbio.2011.09.006]
- 27 **Yilmaz Y**. Review article: fructose in non-alcoholic fatty liver disease. *Aliment Pharmacol Ther* 2012; **35**: 1135-1144 [PMID: 22469071 DOI: 10.1111/j.1365-2036.2012.05080.x]
- 28 **Aeberli I**, Zimmermann MB, Molinari L, Lehmann R, I' Allemand D, Spinass GA, Berneis K. Fructose intake is a predictor of LDL particle size in overweight schoolchildren. *Am J Clin Nutr* 2007; **86**: 1174-1178 [PMID: 17921399]
- 29 **Chiu S**, Sievenpiper JL, de Souza RJ, Cozma AI, Mirrahimi A, Carleton AJ, Ha V, Di Buono M, Jenkins AL, Leiter LA, Wolever TM, Don-Wauchope AC, Beyene J, Kendall CW, Jenkins DJ. Effect of fructose on markers of non-alcoholic fatty liver disease (NAFLD): a systematic review and meta-analysis of controlled feeding trials. *Eur J Clin Nutr* 2014; **68**: 416-423 [PMID: 24569542 DOI: 10.1038/ejcn.2014.8]
- 30 **Hao W**, Wong OY, Liu X, Lee P, Chen Y, Wong KK. ω -3 fatty acids suppress inflammatory cytokine production by macrophages and hepatocytes. *J Pediatr Surg* 2010; **45**: 2412-2418 [PMID: 21129557 DOI: 10.1016/j.jpedsurg.2010.08.044]
- 31 **Ronis MJ**, Korourian S, Zipperman M, Hakkak R, Badger TM. Dietary saturated fat reduces alcoholic hepatotoxicity in rats by altering fatty acid metabolism and membrane composition. *J Nutr* 2004; **134**: 904-912 [PMID: 15051845]

- 32 **Kennedy A**, Martinez K, Chuang CC, LaPoint K, McIntosh M. Saturated fatty acid-mediated inflammation and insulin resistance in adipose tissue: mechanisms of action and implications. *J Nutr* 2009; **139**: 1-4 [PMID: 19056664 DOI: 10.3945/jn.108.098269]
- 33 **Oller do Nascimento CM**, Ribeiro EB, Oyama LM. Metabolism and secretory function of white adipose tissue: effect of dietary fat. *An Acad Bras Cienc* 2009; **81**: 453-466 [PMID: 19722015]
- 34 **Jaeschke H**. Reactive oxygen and mechanisms of inflammatory liver injury: Present concepts. *J Gastroenterol Hepatol* 2011; **26 Suppl 1**: 173-179 [PMID: 21199529 DOI: 10.1111/j.1440-1746.2010.06592.x]
- 35 **Ibrahim SH**, Kohli R, Gores GJ. Mechanisms of lipotoxicity in NAFLD and clinical implications. *J Pediatr Gastroenterol Nutr* 2011; **53**: 131-140 [PMID: 21629127 DOI: 10.1097/MPG.0b013e31822578db]
- 36 **McGarry JD**, Foster DW. Regulation of hepatic fatty acid oxidation and ketone body production. *Annu Rev Biochem* 1980; **49**: 395-420 [PMID: 6157353]
- 37 **Chahal N**, Manhiot C, Wong H, McCrindle BW. Effectiveness of Omega-3 Polysaturated Fatty Acids (Fish Oil) Supplementation for Treating Hypertriglyceridemia in Children and Adolescents. *Clin Pediatr (Phila)* 2014; **53**: 645-651 [PMID: 24647701]
- 38 **Federation of American Societies for Experimental Biology**. Anti-inflammatory effects of omega 3 fatty acid in fish oil linked to lowering of prostaglandin. Rockville: Science Daily, 2006
- 39 **Dieter P**, Scheibe R, Bezugla Y, Matthé E, Schuch S, Treffkorn L, Bernard B, Kamionka S, Kolada A. The regulatory role of prostaglandin E2 in liver (patho) physiology is controlled at its site of synthesis and its action on the receptors. *Comp Hepatol* 2004; **3 Suppl 1**: S35 [PMID: 14960187]
- 40 **Hossain MA**, Wakabayashi H, Izuishi K, Okano K, Yachida S, Maeta H. The role of prostaglandins in liver ischemia-reperfusion injury. *Curr Pharm Des* 2006; **12**: 2935-2951 [PMID: 16918423]
- 41 **Kawahara K**, Hohjoh H, Inazumi T, Tsuchiya S, Sugimoto Y. Prostaglandin E2-induced inflammation: Relevance of prostaglandin E receptors. *Biochim Biophys Acta* 2014; Epub ahead of print [PMID: 25038274 DOI: 10.1016/j.bbailp.2014.07.008]
- 42 **Mater MK**, Thelen AP, Jump DB. Arachidonic acid and PGE2 regulation of hepatic lipogenic gene expression. *J Lipid Res* 1999; **40**: 1045-1052 [PMID: 10357836]
- 43 **Serhan CN**. Novel omega -- 3-derived local mediators in anti-inflammation and resolution. *Pharmacol Ther* 2005; **105**: 7-21 [PMID: 15626453]
- 44 **Takahashi M**, Tsuboyama-Kasaoka N, Nakatani T, Ishii M, Tsutsumi S, Aburatani H, Ezaki O. Fish oil feeding alters liver gene expressions to defend against PPARalpha activation and ROS production. *Am J Physiol Gastrointest Liver Physiol* 2002; **282**: G338-G348 [PMID: 11804856]
- 45 **St-Jules DE**, Watters CA, Brunt EM, Wilkens LR, Novotny R, Belt P, Lavine JE. Estimation of fish and ω -3 fatty acid intake in pediatric nonalcoholic fatty liver disease. *J Pediatr Gastroenterol Nutr* 2013; **57**: 627-633 [PMID: 24177784 DOI: 10.1097/MPG.0b013e3182a1df77]
- 46 **Chirico V**, Lacquaniti A, Salpietro V, Munafò C, Calabrò MP, Buemi M, Arrigo T, Salpietro C. High-mobility group box 1 (HMGB1) in childhood: from bench to bedside. *Eur J Pediatr* 2014; **173**: 1123-1136 [PMID: 24809802 DOI: 10.1007/s00431-014-2327-1]
- 47 **Ashton Acton Q**. Eating Disorders, Nutrition, and Digestive Medicine: 2013 Edition. Available from: URL: <http://books.google.it/books?isbn=1490111816>
- 48 **Arrigo T**, Chirico V, Salpietro V, Munafò C, Ferrau V, Gitto E, Lacquaniti A, Salpietro C. High-mobility group protein B1: a new biomarker of metabolic syndrome in obese children. *Eur J Endocrinol* 2013; **168**: 631-638 [PMID: 23384711 DOI: 10.1530/EJE-13-0037]
- 49 **Lund AS**, Hasselbalch AL, Gamborg M, Skogstrand K, Hougaard DM, Heitmann BL, Kyvik KO, Sørensen TI, Jess T. N-3 polyunsaturated fatty acids, body fat and inflammation. *Obes Facts* 2013; **6**: 369-379 [PMID: 23970146 DOI: 10.1159/000354663]
- 50 **Novak EM**, Keller BO, Innis SM. Metabolic development in the liver and the implications of the n-3 fatty acid supply. *Am J Physiol Gastrointest Liver Physiol* 2012; **302**: G250-G259 [PMID: 22094600 DOI: 10.1152/ajpgi.00189.2011]
- 51 **Popescu LA**, Virgolici B, Lixandru D, Miricescu D, Condrut E, Timnea O, Ranetti AE, Militaru M, Mohora M, Zăgrea L. Effect of diet and omega-3 fatty acids in NAFLD. *Rom J Morphol Embryol* 2013; **54**: 785-790 [PMID: 24322028]
- 52 **Janczyk W**, Socha P, Lebensztejn D, Wierzbicka A, Mazur A, Neuhoff-Murawska J, Matusik P. Omega-3 fatty acids for treatment of non-alcoholic fatty liver disease: design and rationale of randomized controlled trial. *BMC Pediatr* 2013; **13**: 85 [PMID: 23702094 DOI: 10.1186/1471-2431-13-85]
- 53 **Giraldo Villa A**, Henao Roldan C, García Loboguerrero F, Martínez Volkmar MI, Contreras Ramírez MM, Ruiz Navas P. [Use of fish oil lipid emulsions in hospitalized patients under 18 years old with abnormal results in liver tests associated with total parental nutrition]. *Nutr Hosp* 2014; **29**: 844-851 [PMID: 24679026 DOI: 10.3305/nh.2014.29.4.7209]
- 54 **Lee JH**. Polyunsaturated Fatty acids in children. *Pediatr Gastroenterol Hepatol Nutr* 2013; **16**: 153-161 [PMID: 24224148 DOI: 10.5223/pghn.2013.16.3.153]
- 55 **Traber MG**, Stevens JF. Vitamins C and E: beneficial effects from a mechanistic perspective. *Free Radic Biol Med* 2011; **51**: 1000-1013 [PMID: 21664268 DOI: 10.1016/j.freeradbiomed.2011.05.017]
- 56 **Traber MG**, Atkinson J. Vitamin E, antioxidant and nothing more. *Free Radic Biol Med* 2007; **43**: 4-15 [PMID: 17561088]
- 57 **Shaik-Dasthagirisahab YB**, Varvara G, Murmura G, Saggini A, Caraffa A, Antinolfi P, Tete' S, Tripodi D, Conti F, Cianchetti E, Toniato E, Rosati M, Speranza L, Pantalone A, Saggini R, Tei M, Speziali A, Conti P, Theoharides TC, Pandolfi F. Role of vitamins D, E and C in immunity and inflammation. *J Biol Regul Homeost Agents* 2013; **27**: 291-295 [PMID: 23830380]
- 58 **Carr AC**, Zhu BZ, Frei B. Potential antiatherogenic mechanisms of ascorbate (vitamin C) and alpha-tocopherol (vitamin E). *Circ Res* 2000; **87**: 349-354 [PMID: 10969031]
- 59 **Kawanaka M**, Mahmood S, Niiyama G, Izumi A, Kamei A, Ikeda H, Suehiro M, Togawa K, Sasagawa T, Okita M, Nakamura H, Yodoi J, Yamada G. Control of oxidative stress and reduction in biochemical markers by Vitamin E treatment in patients with nonalcoholic steatohepatitis: a pilot study. *Hepatol Res* 2004; **29**: 39-41 [PMID: 15135345]
- 60 **Nobili V**, Manco M, Devito R, Ciampalini P, Piemonte F, Marcellini M. Effect of vitamin E on aminotransferase levels and insulin resistance in children with non-alcoholic fatty liver disease. *Aliment Pharmacol Ther* 2006; **24**: 1553-1561 [PMID: 17206944]
- 61 **Lavine JE**. Vitamin E treatment of nonalcoholic steatohepatitis in children: a pilot study. *J Pediatr* 2000; **136**: 734-738 [PMID: 10839868]
- 62 **Hasegawa T**, Yoneda M, Nakamura K, Makino I, Terano A. Plasma transforming growth factor-beta1 level and efficacy of alpha-tocopherol in patients with non-alcoholic steatohepatitis: a pilot study. *Aliment Pharmacol Ther* 2001; **15**: 1667-1672 [PMID: 11564008]
- 63 **Sanyal AJ**, Chalasani N, Kowdley KV, McCullough A, Diehl AM, Bass NM, Neuschwander-Tetri BA, Lavine JE, Tonascia J, Unalp A, Van Natta M, Clark J, Brunt EM, Kleiner DE, Hoofnagle JH, Robuck PR. Pioglitazone, vitamin E, or placebo for nonalcoholic steatohepatitis. *N Engl J Med* 2010; **362**: 1675-1685 [PMID: 20427778 DOI: 10.1056/NEJMoa0905556]

- 10.1056/NEJMoa0907929]
- 64 **Lavine JE**, Schwimmer JB, Van Natta ML, Molleston JP, Murray KF, Rosenthal P, Abrams SH, Scheimann AO, Sanyal AJ, Chalasani N, Tonascia J, Ünalp A, Clark JM, Brunt EM, Kleiner DE, Hoofnagle JH, Robuck PR. Effect of vitamin E or metformin for treatment of nonalcoholic fatty liver disease in children and adolescents: the TONIC randomized controlled trial. *JAMA* 2011; **305**: 1659-1668 [PMID: 21521847 DOI: 10.1001/jama.2011.520]
- 65 **Kuhad A**, Chopra K. Attenuation of diabetic nephropathy by tocotrienol: involvement of NFκB signaling pathway. *Life Sci* 2009; **84**: 296-301 [PMID: 19162042 DOI: 10.1016/j.lfs.2008.12.014]
- 66 **Bugianesi E**, Gentilecore E, Manini R, Natale S, Vanni E, Villanova N, David E, Rizzetto M, Marchesini G. A randomized controlled trial of metformin versus vitamin E or prescriptive diet in nonalcoholic fatty liver disease. *Am J Gastroenterol* 2005; **100**: 1082-1090 [PMID: 15842582]
- 67 **Halliwell B**, Gutteridge JM. *Free Radicals in Biology and Medicine*. New York: Oxford University Press, 1999
- 68 **Schultz I**, McKay CM, Graepel R, Phan VC, Wang A, Söderholm J, McKay DM. Indomethacin-induced translocation of bacteria across enteric epithelia is reactive oxygen species-dependent and reduced by vitamin C. *Am J Physiol Gastrointest Liver Physiol* 2012; **303**: G536-G545 [PMID: 22700821 DOI: 10.1152/ajpgi.00125.2012]
- 69 **Wang Y**, Russo TA, Kwon O, Chanock S, Rumsey SC, Levine M. Ascorbate recycling in human neutrophils: induction by bacteria. *Proc Natl Acad Sci USA* 1997; **94**: 13816-13819 [PMID: 9391110]
- 70 **Abhilash PA**, Harikrishnan R, Indira M. Ascorbic acid supplementation down-regulates the alcohol induced oxidative stress, hepatic stellate cell activation, cytotoxicity and mRNA levels of selected fibrotic genes in guinea pigs. *Free Radic Res* 2012; **46**: 204-213 [PMID: 22149461 DOI: 10.3109/10715762.2011.647691]
- 71 **Lirussi F**, Azzalini L, Orlando S, Orlando R, Angelico F. Antioxidant supplements for non-alcoholic fatty liver disease and/or steatohepatitis. *Cochrane Database Syst Rev* 2007; (1): CD004996 [PMID: 17253535]
- 72 **Vos MB**, Colvin R, Belt P, Molleston JP, Murray KF, Rosenthal P, Schwimmer JB, Tonascia J, Unalp A, Lavine JE. Correlation of vitamin E, uric acid, and diet composition with histologic features of pediatric NAFLD. *J Pediatr Gastroenterol Nutr* 2012; **54**: 90-96 [PMID: 22197855 DOI: 10.1097/MPG.0b013e318229da1a]
- 73 **Padayatty SJ**, Katz A, Wang Y, Eck P, Kwon O, Lee JH, Chen S, Corpe C, Dutta A, Dutta SK, Levine M. Vitamin C as an antioxidant: evaluation of its role in disease prevention. *J Am Coll Nutr* 2003; **22**: 18-35 [PMID: 12569111]
- 74 **Himmerich H**, Angheliescu I, Klawe C, Szegedi A. Vitamin B12 and hepatic enzyme serum levels correlate in male alcohol-dependent patients. *Alcohol Alcohol* 2001; **36**: 26-28 [PMID: 11139412]
- 75 **Guéant JL**, Caillerez-Fofou M, Battaglia-Hsu S, Alberto JM, Freund JN, Dulluc I, Adjalla C, Maury F, Merle C, Nicolas JP, Namour F, Daval JL. Molecular and cellular effects of vitamin B12 in brain, myocardium and liver through its role as co-factor of methionine synthase. *Biochimie* 2013; **95**: 1033-1040 [PMID: 23415654 DOI: 10.1016/j.biochi.2013.01.020]
- 76 **Xia YM**, Hill KE, Burk RF. Effect of selenium deficiency on hydroperoxide-induced glutathione release from the isolated perfused rat heart. *J Nutr* 1985; **115**: 733-742 [PMID: 3998867]
- 77 **Loguercio C**, De Girolamo V, Federico A, Feng SL, Crafa E, Cataldi V, Gialanella G, Moro R, Del Vecchio Blanco C. Relationship of blood trace elements to liver damage, nutritional status, and oxidative stress in chronic nonalcoholic liver disease. *Biol Trace Elem Res* 2001; **81**: 245-254 [PMID: 11575681]
- 78 **Rayssiguier Y**, Durlach J, Gueux E, Rock E, Mazur A. Magnesium and ageing. I. Experimental data: importance of oxidative damage. *Magnes Res* 1993; **6**: 369-378 [PMID: 8155489]
- 79 **Keen CL**, Ensunsa JL, Watson MH, Baly DL, Donovan SM, Monaco MH, Clegg MS. Nutritional aspects of manganese from experimental studies. *Neurotoxicology* 1999; **20**: 213-223 [PMID: 10385885]
- 80 **Aruoma OI**. Free radicals, oxidative stress, and antioxidants in human health and disease. *J Am Chem Soc* 1998; **75**: 199
- 81 **Hosseini MJ**, Shaki F, Ghazi-Khansari M, Pourahmad J. Toxicity of copper on isolated liver mitochondria: impairment at complexes I, II, and IV leads to increased ROS production. *Cell Biochem Biophys* 2014; **70**: 367-381 [PMID: 24691927 DOI: 10.1007/s12013-014-9922-7]
- 82 **Seth R**, Yang S, Choi S, Sabeen M, Roberts EA. In vitro assessment of copper-induced toxicity in the human hepatoma line, Hep G2. *Toxicol In Vitro* 2004; **18**: 501-509 [PMID: 15130608]
- 83 **Lang PA**, Schenck M, Nicolay JP, Becker JU, Kempe DS, Lupescu A, Koka S, Eisele K, Klarl BA, Rübber H, Schmid KW, Mann K, Hildenbrand S, Hefter H, Huber SM, Wieder T, Erhardt A, Häussinger D, Gulbins E, Lang F. Liver cell death and anemia in Wilson disease involve acid sphingomyelinase and ceramide. *Nat Med* 2007; **13**: 164-170 [PMID: 17259995]
- 84 **Roberts EA**, Sarkar B. Liver as a key organ in the supply, storage, and excretion of copper. *Am J Clin Nutr* 2008; **88**: 851S-854S [PMID: 18779307]
- 85 **Cousins RJ**. Toward a molecular understanding of zinc metabolism. *Clin Physiol Biochem* 1986; **4**: 20-30 [PMID: 2420502]
- 86 **Young IS**, Trouton TG, Torney JJ, McMaster D, Callender ME, Trimble ER. Antioxidant status and lipid peroxidation in hereditary haemochromatosis. *Free Radic Biol Med* 1994; **16**: 393-397 [PMID: 8063202]
- 87 **Bartal M**, Mazor D, Dvilansky A, Meyerstein N. Iron deficiency anemia: recovery from in vitro oxidative stress. *Acta Haematol* 1993; **90**: 94-98 [PMID: 8285025]
- 88 **Ferro E**, Visalli G, Civa R, La Rosa MA, Randazzo Papa G, Baluce B, D'Ascola DG, Piraino B, Salpietro C, Di Pietro A. Oxidative damage and genotoxicity biomarkers in transfused and untransfused thalassemic subjects. *Free Radic Biol Med* 2012; **53**: 1829-1837 [PMID: 22995637 DOI: 10.1016/j.freeradbiomed.2012.08.592]
- 89 **Evans P**. Oxidative damage and genotoxicity biomarkers in transfused and untransfused thalassemic subjects. Micronutrients: oxidant/antioxidant status. *Br J Nutr* 2001; **85**: 67-74
- 90 **Mantena SK**, King AL, Andringa KK, Eccleston HB, Bailey SM. Mitochondrial dysfunction and oxidative stress in the pathogenesis of alcohol- and obesity-induced fatty liver diseases. *Free Radic Biol Med* 2008; **44**: 1259-1272 [PMID: 18242193 DOI: 10.1016/j.freeradbiomed.2007.12.029]
- 91 **Kowdley KV**, Belt P, Wilson LA, Yeh MM, Neuschwander-Tetri BA, Chalasani N, Sanyal AJ, Nelson JE. Serum ferritin is an independent predictor of histologic severity and advanced fibrosis in patients with nonalcoholic fatty liver disease. *Hepatology* 2012; **55**: 77-85 [PMID: 21953442 DOI: 10.1002/hep.24706]
- 92 **Aigner E**, Theurl I, Theurl M, Lederer D, Haufe H, Dietze O, Strasser M, Datz C, Weiss G. Pathways underlying iron accumulation in human nonalcoholic fatty liver disease. *Am J Clin Nutr* 2008; **87**: 1374-1383 [PMID: 18469261]
- 93 **Ahmed U**, Latham PS, Oates PS. Interactions between hepatic iron and lipid metabolism with possible relevance to steatohepatitis. *World J Gastroenterol* 2012; **18**: 4651-4658 [PMID: 23002334]
- 94 **Ganne-Carrié N**, Christidis C, Chastang C, Ziolk M, Chapel F, Imbert-Bismut F, Trinchet JC, Guettier C, Beaugrand M.

- Liver iron is predictive of death in alcoholic cirrhosis: a multivariate study of 229 consecutive patients with alcoholic and/or hepatitis C virus cirrhosis: a prospective follow up study. *Gut* 2000; **46**: 277-282 [PMID: 10644325]
- 95 **Mueller S**, Millonig G, Seitz HK. Alcoholic liver disease and hepatitis C: a frequently underestimated combination. *World J Gastroenterol* 2009; **15**: 3462-3471 [PMID: 19630099]
- 96 **Girelli D**, Pasino M, Goodnough JB, Nemeth E, Guido M, Castagna A, Busti F, Camprostrini N, Martinelli N, Vantini I, Corrocher R, Ganz T, Fattovich G. Reduced serum hepcidin levels in patients with chronic hepatitis C. *J Hepatol* 2009; **51**: 845-852 [PMID: 19729219 DOI: 10.1016/j.jhep.2009.06.027]
- 97 **Tanaka H**, Fujita N, Sugimoto R, Urawa N, Horriike S, Kobayashi Y, Iwasa M, Ma N, Kawanishi S, Watanabe S, Kaito M, Takei Y. Hepatic oxidative DNA damage is associated with increased risk for hepatocellular carcinoma in chronic hepatitis C. *Br J Cancer* 2008; **98**: 580-586 [PMID: 18231107 DOI: 10.1038/sj.bjc.6604204]
- 98 **Food and Agriculture Organization of the United Nations**. Guidelines for the Evaluation of probiotics in Food, 2002. Available from: URL: ftp://ftp.fao.org/es/esn/food/wgreport2.pdf
- 99 **Takaki A**, Kawai D, Yamamoto K. Multiple hits, including oxidative stress, as pathogenesis and treatment target in non-alcoholic steatohepatitis (NASH). *Int J Mol Sci* 2013; **14**: 20704-20728 [PMID: 24132155 DOI: 10.3390/ijms141020704]
- 100 **Roh YS**, Seki E. Toll-like receptors in alcoholic liver disease, non-alcoholic steatohepatitis and carcinogenesis. *J Gastroenterol Hepatol* 2013; **28** Suppl 1: 38-42 [PMID: 23855294 DOI: 10.1111/jgh.12019]
- 101 **Arora S**, Kaur IP, Chopra K, Rishi P. Efficiency of double layered microencapsulated probiotic to modulate proinflammatory molecular markers for the management of alcoholic liver disease. *Mediators Inflamm* 2014; **2014**: 715130 [PMID: 24966470 DOI: 10.1155/2014/715130]
- 102 **Zhou YK**, Qin HL, Zhang M, Shen TY, Chen HQ, Ma YL, Chu ZX, Zhang P, Liu ZH. Effects of *Lactobacillus plantarum* on gut barrier function in experimental obstructive jaundice. *World J Gastroenterol* 2012; **18**: 3977-3991 [PMID: 22912548 DOI: 10.3748/wjg.v18.i30.3977]
- 103 **Reid G**, Jass J, Sebulsy MT, McCormick JK. Potential uses of probiotics in clinical practice. *Clin Microbiol Rev* 2003; **16**: 658-672 [PMID: 14557292]
- 104 **Lionetti E**, Francavilla R, Castellazzi AM, Arrigo T, Labò E, Leonardi S, Ciprandi G, Miraglia Del Giudice M, Salpietro V, Salpietro C, La Rosa M. Probiotics and *Helicobacter pylori* infection in children. *J Biol Regul Homeost Agents* 2012; **26**: S69-S76 [PMID: 22691253]
- 105 **del Giudice MM**, Leonardi S, Ciprandi G, Galdo F, Gubitosi A, La Rosa M, Salpietro C, Marseglia G, Perrone L. Probiotics in childhood: allergic illness and respiratory infections. *J Clin Gastroenterol* 2012; **46** Suppl: S69-S72 [PMID: 22955363 DOI: 10.1097/MCG.0b013e318266fea7]
- 106 **Salpietro C**, Rigoli L, Miraglia Del Giudice M, Cuppari C, Di Bella C, Salpietro A, Maiello N, La Rosa M, Marseglia GL, Leonardi S, Briuglia S, Ciprandi G. TLR2 and TLR4 gene polymorphisms and atopic dermatitis in Italian children: a multicenter study. *Int J Immunopathol Pharmacol* 2011; **24**: 33-40 [PMID: 22032785]
- 107 **Rigoli L**, Briuglia S, Caimmi S, Ferrau V, Gallizzi R, Leonardi S, La Rosa M, Salpietro C. Gene-environment interaction in childhood asthma. *Int J Immunopathol Pharmacol* 2011; **24**: 41-47 [PMID: 22032786]
- 108 **Mencarelli A**, Cipriani S, Renga B, Bruno A, D'Amore C, Distrutti E, Fiorucci S. VSL#3 resets insulin signaling and protects against NASH and atherosclerosis in a model of genetic dyslipidemia and intestinal inflammation. *PLoS One* 2012; **7**: e45425 [PMID: 23029000 DOI: 10.1371/journal.pone.0045425]
- 109 **Rashid SK**, Khodja NI, Auger C, Alhosin M, Boehm N, Oswald-Mammosser M, Schini-Kerth VB. Probiotics (VSL#3) prevent endothelial dysfunction in rats with portal hypertension: role of the angiotensin system. *PLoS One* 2014; **9**: e97458 [PMID: 24832090 DOI: 10.1371/journal.pone.0097458]
- 110 **Lirussi F**, Mastropasqua E, Orando S, Orlando R. Probiotics for non-alcoholic fatty liver disease and/or steatohepatitis. *Cochrane Database Syst Rev* 2007; (1): CD005165 [PMID: 17253543]
- 111 **Aller R**, De Luis DA, Izaola O, Conde R, Gonzalez Sagrado M, Primo D, De La Fuente B, Gonzalez J. Effect of a probiotic on liver aminotransferases in nonalcoholic fatty liver disease patients: a double blind randomized clinical trial. *Eur Rev Med Pharmacol Sci* 2011; **15**: 1090-1095 [PMID: 22013734]
- 112 **Vajro P**, Mandato C, Licenziati MR, Franzese A, Vitale DF, Lenta S, Caropreso M, Vallone G, Meli R. Effects of *Lactobacillus rhamnosus* strain GG in pediatric obesity-related liver disease. *J Pediatr Gastroenterol Nutr* 2011; **52**: 740-743 [PMID: 21505361 DOI: 10.1097/MPG.0b013e31821f9b85]
- 113 **Kirpich IA**, Solovieva NV, Leikhter SN, Shidakova NA, Lebedeva OV, Sidorov PI, Bazhukova TA, Soloviev AG, Barve SS, McClain CJ, Cave M. Probiotics restore bowel flora and improve liver enzymes in human alcohol-induced liver injury: a pilot study. *Alcohol* 2008; **42**: 675-682 [PMID: 19038698 DOI: 10.1016/j.alcohol.2008.08.006]
- 114 **Blaszuk H**, Wild PJ, Oliveira A, Kelly DG, Burgart LJ. Hepatic copper in patients receiving long-term total parenteral nutrition. *J Clin Gastroenterol* 2005; **39**: 318-320 [PMID: 15758626]
- 115 **Zambrano E**, El-Hennawy M, Ehrenkranz RA, Zelterman D, Reyes-Múgica M. Total parenteral nutrition induced liver pathology: an autopsy series of 24 newborn cases. *Pediatr Dev Pathol* 2004; **7**: 425-432 [PMID: 15547767]
- 116 **Tulikoura I**, Vuori E. Effect of total parenteral nutrition on the zinc, copper, and manganese status of patients with catabolic disease. *Scand J Gastroenterol* 1986; **21**: 421-427 [PMID: 3088719]
- 117 **Hamacher S**, Matern S, Roeb E. [Extracellular matrix -- from basic research to clinical significance. An overview with special consideration of matrix metalloproteinases]. *Dtsch Med Wochenschr* 2004; **129**: 1976-1980 [PMID: 15375740]
- 118 **Tazuke Y**, Drongowski RA, Btaiche I, Coran AG, Teitelbaum DH. Effects of lipid administration on liver apoptotic signals in a mouse model of total parenteral nutrition (TPN). *Pediatr Surg Int* 2004; **20**: 224-228 [PMID: 15034728]
- 119 **Ferri KF**, Kroemer G. Organelle-specific initiation of cell death pathways. *Nat Cell Biol* 2001; **3**: E255-E263 [PMID: 11715037]
- 120 **Leist M**, Single B, Castoldi AF, Kühnle S, Nicotera P. Intracellular adenosine triphosphate (ATP) concentration: a switch in the decision between apoptosis and necrosis. *J Exp Med* 1997; **185**: 1481-1486 [PMID: 9126928]
- 121 **Halliwell B**, Chirico S. Lipid peroxidation: its mechanism, measurement, and significance. *Am J Clin Nutr* 1993; **57**: 715S-724S; discussion 724S-725S [PMID: 8475889]
- 122 **Pierro A**, van Saene HK, Donnell SC, Hughes J, Ewan C, Nunn AJ, Lloyd DA. Microbial translocation in neonates and infants receiving long-term parenteral nutrition. *Arch Surg* 1996; **131**: 176-179 [PMID: 8611075]
- 123 **Gore DC**, Chinkes D, Heggers J, Herndon DN, Wolf SE, Desai M. Association of hyperglycemia with increased mortality after severe burn injury. *J Trauma* 2001; **51**: 540-544 [PMID: 11535907]
- 124 **Barber AE**, Jones WG, Minei JP, Fahey TJ, Lowry SF, Shires GT. Bacterial overgrowth and intestinal atrophy in the etiology of gut barrier failure in the rat. *Am J Surg* 1991; **161**: 300-304 [PMID: 1990885]
- 125 **Bhatia J**, Moslen MT, Haque AK, McCleery R, Rassin DK. Total parenteral nutrition-associated alterations in hepa-

- tobiliary function and histology in rats: is light exposure a clue? *Pediatr Res* 1993; **33**: 487-492 [PMID: 8511021]
- 126 **Schepens MA**, Roelofs HM, Peters WH, Wanten GJ. No evidence for oxidative stress in patients on home parenteral nutrition. *Clin Nutr* 2006; **25**: 939-948 [PMID: 16777272]
- 127 **Moosivand A**, Abrishami R, Abdollahi M, Ahmadi A, Mojtahedzadeh M. The incidences of oxidative-stress occurrence following two metabolic support measures in critically ill patients. *J Pharm Care* 2013; **1**: 3-7
- 128 **Miloudi K**, Comte B, Rouleau T, Montoudis A, Levy E, Lavoie JC. The mode of administration of total parenteral nutrition and nature of lipid content influence the generation of peroxides and aldehydes. *Clin Nutr* 2012; **31**: 526-534 [PMID: 22230256 DOI: 10.1016/j.clnu.2011.12.012]

P- Reviewer: El-Shabrawi MH, Miyoshi E **S- Editor:** Ma YJ
L- Editor: Stewart GJ **E- Editor:** Ma S





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045